



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p><b>(51) International Patent Classification <sup>6</sup> :</b>  <b>C07D 231/38, A61K 31/415, C07D 401/10, 403/12, 401/12, A61K 31/44</b></p>	<p><b>A1</b></p>	<p><b>(11) International Publication Number:</b> <b>WO 99/57101</b></p> <p><b>(43) International Publication Date:</b> 11 November 1999 (11.11.99)</p>		
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top; padding: 5px;"> <p><b>(21) International Application Number:</b> PCT/EP99/02879</p> <p><b>(22) International Filing Date:</b> 28 April 1999 (28.04.99)</p> <p><b>(30) Priority Data:</b>  60/084,250                      5 May 1998 (05.05.98)                      US  60/122,410                      2 March 1999 (02.03.99)                      US</p> <p><b>(71) Applicant:</b> F. HOFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH).</p> <p><b>(72) Inventors:</b> LABADIE, Sharada, Shenvi; 1618 Kamsack Drive, Sunnyvale, CA 94087 (US). ROTSTEIN, David, Mark; 939 Lome Way, Sunnyvale, CA 94087 (US). SJOGREN, Eric, Brian; 442 Dell Avenue, Mountain View, CA 94043 (US). TALAMAS, Francisco, Xavier; 1369 Geneva Street, San Carlos, CA 94070 (US).</p> <p><b>(74) Agent:</b> LÖSCHNER, Thomas; Grenzacherstrasse 124, CH-4070 Basle (CH).</p> </td> <td style="width: 50%; vertical-align: top; padding: 5px;"> <p><b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b>  <i>With international search report.  Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> </td> </tr> </table>			<p><b>(21) International Application Number:</b> PCT/EP99/02879</p> <p><b>(22) International Filing Date:</b> 28 April 1999 (28.04.99)</p> <p><b>(30) Priority Data:</b>  60/084,250                      5 May 1998 (05.05.98)                      US  60/122,410                      2 March 1999 (02.03.99)                      US</p> <p><b>(71) Applicant:</b> F. HOFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH).</p> <p><b>(72) Inventors:</b> LABADIE, Sharada, Shenvi; 1618 Kamsack Drive, Sunnyvale, CA 94087 (US). ROTSTEIN, David, Mark; 939 Lome Way, Sunnyvale, CA 94087 (US). SJOGREN, Eric, Brian; 442 Dell Avenue, Mountain View, CA 94043 (US). TALAMAS, Francisco, Xavier; 1369 Geneva Street, San Carlos, CA 94070 (US).</p> <p><b>(74) Agent:</b> LÖSCHNER, Thomas; Grenzacherstrasse 124, CH-4070 Basle (CH).</p>	<p><b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b>  <i>With international search report.  Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p><b>(21) International Application Number:</b> PCT/EP99/02879</p> <p><b>(22) International Filing Date:</b> 28 April 1999 (28.04.99)</p> <p><b>(30) Priority Data:</b>  60/084,250                      5 May 1998 (05.05.98)                      US  60/122,410                      2 March 1999 (02.03.99)                      US</p> <p><b>(71) Applicant:</b> F. HOFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH).</p> <p><b>(72) Inventors:</b> LABADIE, Sharada, Shenvi; 1618 Kamsack Drive, Sunnyvale, CA 94087 (US). ROTSTEIN, David, Mark; 939 Lome Way, Sunnyvale, CA 94087 (US). SJOGREN, Eric, Brian; 442 Dell Avenue, Mountain View, CA 94043 (US). TALAMAS, Francisco, Xavier; 1369 Geneva Street, San Carlos, CA 94070 (US).</p> <p><b>(74) Agent:</b> LÖSCHNER, Thomas; Grenzacherstrasse 124, CH-4070 Basle (CH).</p>	<p><b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b>  <i>With international search report.  Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>			
<p><b>(54) Title:</b> PYRAZOLE DERIVATIVES AS P-38 MAP KINASE INHIBITORS</p> <div style="text-align: center; margin: 20px 0;"> <div style="position: absolute; left: 600px; top: 50px;">(I)</div> </div>				
<p><b>(57) Abstract</b></p> <p>The present invention relates to certain pyrazole derivatives of Formula (I) that are p-38 MAP kinase inhibitors, pharmaceutical compositions containing them, methods for their use, and methods for preparing these compounds.</p>				

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## PYRAZOLE DERIVATIVES AS P-38 MAP KINASE INHIBITORS

This invention relates to certain pyrazole derivatives that inhibit p38 MAP kinase, pharmaceutical compositions containing them, their use, intermediates and processes for preparing these compounds.

5           TNF and IL-1 have been shown to be central players in the pathological processes underlying many chronic inflammatory and autoimmune diseases. IL-1 is implicated in mediating or exacerbating diseases such as rheumatoid arthritis ((*see.*, Arend, W. P. *Arthritis & Rheumatism* **38**(2): 151-160, (1995)), osteoarthritis, bone resorption, toxic  
10 shock syndrome, tuberculosis, atherosclerosis, diabetes, Hodgkin's disease (*see.*, Benharroch, D.; et. al. *Euro. Cytokine Network* **7**(1): 51-57) and Alzheimer's disease. Excessive or unregulated TNF production has been implicated in mediating or exacerbating diseases such as rheumatoid arthritis ((*see.*, Maini, R. N.; et. al. *APMIS*. **105**(4): 257-263, (1997);  
15 Feldmann, M., *J. of the Royal College of Physicians of London* **30**(6): 560-570, (1996); Lorenz, H. M.; et. al. *J. of Immunology* **156**(4): 1646-1653, (1996)) osteoarthritis, spondylitis, sepsis, septic shock ((*see.*, Abraham, E.; et. al. *JAMA*. **277**(19):1531-1538, (1997), adult respiratory distress syndrome, asthma ((*see.*, Shah, A.; et. al. *Clin. & Exp. Allergy* 1038-1044,  
20 (1995) and Lassalle, P., et. al. *Clin. & Exp. Immunol.* **94**(1): 105-110, (1993)), bone resorption diseases, fever ((*see.*, Cooper, A. L., et. al. *Am. J. of Physiology* **267**(6 Pt. 2): 1431-1436)), encephalomyelitis, demyelination ((*see.*, Klindert, W. E.; et al. *J. of Neuroimmunol.* **72**(2): 163-168, (1997)) and periodontal diseases.


25           Clinical trials with IL-1 and TNF receptor antagonists have shown that blocking the ability of these cytokines to signal through their receptors leads to significant improvement, in humans, in inflammatory diseases. Therefore, modulation of these inflammatory cytokines is  
30 considered one of the most effective strategies to block chronic inflammation and have positive therapeutic outcomes. It has also been shown that p38 MAP kinase plays an important role in the translational control of TNF and IL-1 and is also involved in the biochemical signaling of

- 2 -

these molecules ((see., Lee, J. C., et al. *Nature*. **372** (6508): 739-46, (1994)). Compounds that bind to p38 MAP are effective in inhibiting bone resorption, inflammation, and other immune and inflammation-based pathologies. The characterization of the p38 MAP kinase and its central role in the biosynthesis of TNF and IL-1 have made this kinase an attractive target for the treatment of diseases mediated by these cytokines.

It would therefore be desirable to provide p38 MAP kinase inhibitors and thereby provide a means of combating diseases mediated by pro-inflammatory cytokines such as TNF and IL-1. This invention fulfills this and related needs.

In a first aspect, this invention provides compounds selected from the group of compounds represented by Formula (I):



wherein:

R<sup>1</sup> is hydrogen, acyl or -P(O)(OH)<sub>2</sub>;

R<sup>2</sup> is hydrogen, halo, alkyl or alkylthio;

A is an aryl, heteroaryl or a heterocyclcyl ring optionally fused to a phenyl ring provided that the heterocyclcyl ring is attached to the carbonyl group via a carbon ring atom;

B is an aryl or heteroaryl ring;

R<sup>3</sup> is selected from the group consisting of:

- (a) amino, alkylamino or dialkylamino;
- (b) acylamino;
- (c) optionally substituted heterocyclcyl;
- (d) optionally substituted aryl or heteroaryl;

- 3 -

- (e) heteroalkyl;  
 (f) heteroalkenyl;  
 (g) heteroalkynyl;  
 (h) heteroalkoxy;  
 5 (i) heteroalkylamino;  
 (j) optionally substituted heterocyclalkyl;  
 (k) optionally substituted heterocyclalkenyl;  
 (l) optionally substituted heterocyclalkynyl;  
 10 (m) optionally substituted heterocyclalkoxy or heterocycloxy;  
 (n) optionally substituted heterocyclalkylamino;  
 (o) optionally substituted heterocyclalkylcarbonyl;  
 (p) heteroalkylcarbonyl;  
 15 (q)  $\text{-NHSO}_2\text{R}^6$  where  $\text{R}^6$  is alkyl, heteroalkyl or optionally substituted heterocyclalkyl;  
 (r)  $\text{-NHSO}_2\text{NR}^7\text{R}^8$  where  $\text{R}^7$  and  $\text{R}^8$  are, independently of each other, hydrogen, alkyl or heteroalkyl;  
 20 (s)  $\text{-Y-(alkylene)-R}^9$  where:  
 Y is a single bond,  $\text{-O-}$ ,  $\text{-NH-}$  or  $\text{-S(O)}_n\text{-}$  (where n is an integer from 0 to 2); and  
 $\text{R}^9$  is cyano, optionally substituted heteroaryl,  $\text{-COOH}$ ,  $\text{-COR}^{10}$ ,  $\text{-COOR}^{11}$ ,  $\text{-CONR}^{12}\text{R}^{13}$ ,  $\text{-SO}_2\text{R}^{14}$ ,  $\text{-SO}_2\text{NR}^{15}\text{R}^{16}$ ,  $\text{-NHSO}_2\text{R}^{17}$  or  $\text{-NHSO}_2\text{NR}^{18}\text{R}^{19}$ , where  
 25  $\text{R}^{10}$  is alkyl or optionally substituted heterocycle,  $\text{R}^{11}$  is alkyl, and  $\text{R}^{12}$ ,  $\text{R}^{13}$ ,  $\text{R}^{14}$ ,  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{17}$ ,  $\text{R}^{18}$  and  $\text{R}^{19}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;  
 30 (t)  $\text{-C(=NR}^{20}\text{)(NR}^{21}\text{R}^{22}\text{)}$  where  $\text{R}^{20}$ ,  $\text{R}^{21}$  and  $\text{R}^{22}$  independently represent hydrogen, alkyl or hydroxy, or  $\text{R}^{20}$  and  $\text{R}^{21}$  together are  $\text{-(CH}_2\text{)}_n\text{-}$  where n is 2 or 3 and  $\text{R}^{22}$  is hydrogen or alkyl;

- 4 -

- (u)  $\text{-NHC(X)NR}^{23}\text{R}^{24}$  where X is -O- or -S-, and  $\text{R}^{23}$  and  $\text{R}^{24}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;
- (v)  $\text{-CONR}^{25}\text{R}^{26}$  where  $\text{R}^{25}$  and  $\text{R}^{26}$  independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclalkyl, or  $\text{R}^{25}$  and  $\text{R}^{26}$  together with the nitrogen to which they are attached form an optionally substituted heterocycl ring;
- (w)  $\text{-S(O)}_n\text{R}^{27}$  where n is an integer from 0 to 2, and  $\text{R}^{27}$  is alkyl, heteroalkyl, optionally substituted heterocyclalkyl or  $\text{-NR}^{28}\text{R}^{29}$  where  $\text{R}^{28}$  and  $\text{R}^{29}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;
- (x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;
- (y) arylaminoalkylene or heteroarylaminomethylene;
- (z) Z-alkylene- $\text{NR}^{30}\text{R}^{31}$  or Z-alkylene- $\text{OR}^{32}$  where Z is -NH-, -N(alkyl)- or -O-, and  $\text{R}^{30}$ ,  $\text{R}^{31}$  and  $\text{R}^{32}$  are independently of each other, hydrogen, alkyl or heteroalkyl;
- (aa)  $\text{-OC(O)-alkylene-CO}_2\text{H}$  or  $\text{-OC(O)-NR'R''}$  (where  $\text{R'}$  and  $\text{R''}$  are independently hydrogen or alkyl); and
- (bb) heteroarylalkenylene or heteroarylalkynylene;
- $\text{R}^4$  is selected from the group consisting of:
- (a) hydrogen;
- (b) halo;
- (c) alkyl;
- (d) alkoxy; and
- (e) hydroxy;
- $\text{R}^5$  is selected from the group consisting of:
- (a) hydrogen;

- 5 -

- (b) halo;
- (c) alkyl;
- (d) haloalkyl;
- (e) thioalkyl;
- 5 (f) hydroxy;
- (g) amino;
- (h) alkylamino;
- (i) dialkylamino;
- (j) heteroalkyl;
- 10 (k) optionally substituted heterocycle;
- (l) optionally substituted heterocyclalkyl;
- (m) optionally substituted heterocyclalkoxy;
- (n) alkylsulfonyl;
- 15 (o) aminosulfonyl, mono-alkylaminosulfonyl or di-alkylaminosulfonyl;
- (p) heteroalkoxy; and
- (q) carboxy;

R<sup>6</sup> is selected from a group consisting of:

- (a) hydrogen;
- 20 (b) halo;
- (c) alkyl; and
- (d) alkoxy;

prodrugs, individual isomers, mixtures of isomers and  
pharmaceutically acceptable salts thereof. In a second aspect, this  
25 invention provides processes for preparing compounds of Formula  
(I). In a third aspect this invention provides intermediates for  
preparing compounds of Formula (I).

In a fourth aspect, this invention provides medicaments or  
30 pharmaceutical compositions containing a therapeutically effective amount  
of a compound of Formula (I) or its pharmaceutically acceptable salt and a  
pharmaceutically acceptable excipient.

- 6 -

In a fifth aspect, this invention relates to the use of a compound of Formula (I).

Unless otherwise stated, the following terms used in the  
5 specification and claims have the meanings given below:

"Alkyl" means a linear saturated monovalent hydrocarbon radical of one to six carbon atoms or a branched saturated monovalent hydrocarbon radical of three to six carbon atoms, e.g., methyl, ethyl, propyl, 2-propyl,  
10 pentyl, and the like.

"Alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, e.g., methylene, ethylene, propylene,  
15 2-methylpropylene, pentylene, and the like.

"Alkenyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenyl, propenyl, and the like.  
20

"Alkenylene" means a linear divalent hydrocarbon radical of two to six carbon atoms or a branched divalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenylene, propenylene, and the like.  
25

"Alkynyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched divalent hydrocarbon radical of three to six carbon atoms, containing at least one triple bond, e.g., ethynyl, propynyl, and the like.  
30

"Alkynylene" means a linear divalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to



- 7 -

six carbon atoms, containing at least one triple bond, e.g., ethynylene, propynylene, and the like.

"Alkoxy" means a radical -OR where R is alkyl as defined above,  
5 e.g., methoxy, ethoxy, propoxy, 2-propoxy, the like.

"Acyl" means a radical -C(O)R where R is alkyl or haloalkyl e.g., acetyl, trifluoroacetyl, and the like.

10 "Acylamino" means a radical -NRC(O)R' where R is hydrogen or alkyl, and R' is alkyl, heteroalkyl or optionally substituted heterocyclalkyl, e.g., acetylamino, 2-amino-2-methylpropionamide, and the like.

"Halo" means fluoro, chloro, bromo, or iodo, preferably fluoro and  
15 chloro.

"Haloalkyl" means alkyl substituted with one or more same or different halo atoms, e.g., -CH<sub>2</sub>Cl, -CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CCl<sub>3</sub>, and the like.

20 "Aryl" means a monovalent monocyclic or bicyclic aromatic hydrocarbon radical of 6 to 10 ring atoms e.g., phenyl, 1-naphthyl, 2-naphthyl, and the like. The aryl ring may optionally be fused to a 5-, 6- or 7- membered monocyclic saturated ring optionally containing 1 or 2 heteroatoms independently selected from oxygen, nitrogen or sulfur, the  
25 remaining ring atoms being C where one or two C atoms are optionally replaced by a carbonyl group. Representative aryl radicals with fused rings include, but are not limited to, 2,3-dihydrobenzo[1,4]dioxan, chroman, isochroman, 2,3-dihydrobenzofuran, 1,3-dihydroisobenzofuran, benzo[1,3]dioxole, 1,2,3,4-tetrahydroisoquinoline, 1,2,3,4-  
30 tetrahydroquinoline, 2,3-dihydro-1H-indole, 2,3-dihydro-1H-isoindole, benzimidazol-2-one, 3H-benzoxazol-2-one, and the like.

"Heteroaryl" means a monovalent monocyclic or bicyclic aromatic radical of 5 to 10 ring atoms containing one, two, or three ring heteroatoms

- 8 -

selected from N, O, or S, the remaining ring atoms being C. The term also includes those radicals where a heteroatom within the ring has been oxidized or quaternized, such as, for example, to form an N-oxide or a quaternary salt. Representative examples include, but are not limited to, thienyl, benzothienyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, quinolinyl, quinoxalinyl, imidazolyl, furanyl, benzofuranyl, thiazolyl, isoxazolyl, benzisoxazolyl, benzimidazolyl, triazolyl, pyrazolyl, pyrrolyl, indolyl, 2-pyridonyl, 4-pyridonyl, N-alkyl-2-pyridonyl, pyrazinonyl, pyridazinonyl, pyrimidinonyl, oxazolonyl, and their corresponding N-oxides, (e.g. pyridyl N-oxide, quinolinyl N-oxide), their quaternary salts and the like.

"Heterocycle" or "heterocyclyl" means a cyclic nonaromatic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from N, O, or S(O)<sub>n</sub> (where n is an integer from 0 to 2), the remaining ring atoms being C where one or two C atoms are optionally replaced by a carbonyl group. The term also includes those radicals where a ring nitrogen atom has been oxidized or quaternized, such as, for example, to form an N-oxide or a quaternary salt. Representative examples include, but are not limited to, tetrahydropyranyl, tetrahydrofuranyl, tetrahydrothiophenyl, piperidino, morpholino, piperazino, pyrrolidino, oxiranyl, dioxane, 1,3-dioxolanyl, 2,2-dimethyl-1,3-dioxalanyl, sulfolanyl, 2-oxazolidonyl, 2-imidazolidonyl, S,S-dioxo-thiomorpholino, and the like.

"Heterocycloamino" means a saturated monovalent cyclic group of 4 to 8 ring atoms, wherein at least one ring atom is N and optionally contains one additional ring atom selected from N or O, the remaining ring atoms being C. The term includes groups such as pyrrolidino, piperidino, morpholino, piperazino and the like.

"Optionally substituted aryl, heteroaryl or heterocyclyl" means an aryl, heteroaryl or heterocyclyl ring as defined above, which is optionally substituted independently with one or two substituents selected from alkyl, phenyl, benzyl, haloalkyl, heteroalkyl, halo, cyano, acyl, -OR (where R is

- 9 -

hydrogen or alkyl), -NRR' (where R and R' are independently selected from hydrogen, alkyl or acyl), -NHCOR (where R is alkyl), -NRS(O)<sub>n</sub>R' (where R is hydrogen or alkyl, n is an integer from 0 to 2 and R' is hydrogen, alkyl or heteroalkyl), -NRS(O)<sub>n</sub>NR'R" (where R is hydrogen or alkyl, n is an integer  
 5 from 0 to 2 and R' and R" are independently hydrogen, alkyl or heteroalkyl), -S(O)<sub>n</sub>R (where n is an integer from 0 to 2 and R is hydrogen, alkyl or heteroalkyl), -S(O)<sub>n</sub>NRR' (where n is an integer from 0 to 2 and R and R' are independently hydrogen, alkyl or heteroalkyl), -COOR, -  
 (alkylene)COOR (where R is hydrogen or alkyl), -CONR'R" or -  
 10 (alkylene)CONR'R" (where R' and R" are independently hydrogen or alkyl).

"Heteroalkyl" means an alkyl radical as defined above, carrying one, two or three substituents selected from -NR<sup>a</sup>R<sup>b</sup>, -OR<sup>c</sup> wherein R<sup>a</sup>, R<sup>b</sup> and R<sup>c</sup> are independently of each other hydrogen, alkyl or acyl, or R<sup>a</sup> and R<sup>b</sup>  
 15 together form heterocycloamino group. Representative examples include, but are not limited to, hydroxymethyl, acetoxymethyl, 3-hydroxypropyl, 1,2-dihydroxyethyl, 2-methoxyethyl, 2-aminoethyl, 2-dimethylaminoethyl, 2-acetylaminomethyl, 3-[pyrrolidin-1-yl]ethyl and the like.

"Heteroalkenyl" means an alkenyl radical as defined above, carrying one or two substituents selected from -NR<sup>a</sup>R<sup>b</sup>, -OR<sup>c</sup> or -S(O)<sub>n</sub>R<sup>d</sup> wherein R<sup>a</sup>, R<sup>b</sup> and R<sup>c</sup> are independently of each other hydrogen or alkyl, and R<sup>d</sup> is alkyl or -NRR' (where R and R' are independently of each other hydrogen or  
 20 alkyl. Representative examples include, but are not limited to, 3-hydroxy-1-propenyl, 3-aminoprop-1-enyl, 2-aminosulfonylethenyl, 2-methylsulfonylethenyl, and the like.

"Heteroalkynyl" means an alkynyl radical as defined above, carrying one or two substituents selected -NR<sup>a</sup>R<sup>b</sup>, -OR<sup>c</sup>, -S(O)<sub>n</sub>R<sup>d</sup> or -S(O)<sub>n</sub>NRR' (where R and R' are independently of each other hydrogen or alkyl)  
 30 wherein R<sup>a</sup>, R<sup>b</sup> and R<sup>c</sup> are independently of each other hydrogen or alkyl, and R<sup>d</sup> is alkyl and n is an integer from zero to two. Representative examples include, but are not limited to, 3-hydroxy-1-propynyl, 3-dimethylaminoprop-1-ynyl and the like.

- 10 -

"Heteroalkoxy" means a radical -OR where R is heteroalkyl group as defined above, e.g., 2-hydroxyethoxy, 3-hydroxypropoxy, 2,3-dihydroxypropoxy, 2-aminoethoxy, and the like.

5

"Heteroalkylamino" means a radical -NR<sup>a</sup>R<sup>b</sup> where R<sup>a</sup> is hydrogen or alkyl, and R<sup>b</sup> is a heteroalkyl group as defined above, e.g., 2-hydroxyethylamino, 3-dimethylaminopropylamino, and the like.

10

" Optionally substituted heterocyclalkyl" means a radical -R<sup>a</sup>R<sup>b</sup> where R<sup>a</sup> is an alkylene group, and R<sup>b</sup> is an optionally substituted heterocycl group as defined above e.g., 2-(morpholin-4-yl)ethyl, 3-(piperidin-1-yl)-2-methylpropyl, and the like.

15

" Optionally substituted heterocyclalkenyl" means a radical -R<sup>a</sup>R<sup>b</sup> where R<sup>a</sup> is an alkenylene group and R<sup>b</sup> is an optionally substituted heterocycl group as defined above e.g., 3-(morpholin-4-yl)prop-1-enyl, 3-(piperidin-1-yl)prop-1-enyl, 3-(4-methylpiperazin-1-yl)prop-1-enyl, and the like.

20

" Optionally substituted heterocyclalkynyl" means a radical -R<sup>a</sup>R<sup>b</sup> where R<sup>a</sup> is an alkynyl group and R<sup>b</sup> is an optionally substituted heterocycl group as defined above e.g., 3-(morpholin-4-yl)prop-1-ynyl, 3-(piperidin-1-yl)prop-1-ynyl, and the like.

25

"Optionally substituted heterocyclalkoxy" means a radical -OR where R is an optionally substituted heterocyclalkyl group as defined above, e.g., 2-(morpholin-4-yl)-ethoxy, 3-(piperazin-1-yl)propoxy, 2-[2-oxopyrrolidin-1-yl]ethoxy, and the like.

30

"Optionally substituted heterocyclalkylamino" means a radical -NR<sup>a</sup>R<sup>b</sup> where R<sup>a</sup> is hydrogen or alkyl and R<sup>b</sup> is an optionally substituted heterocyclalkyl group as defined above, e.g., 2-(pyrrolidin-2-yl)ethylamino, 3-(piperidin-1-yl)propylamino, and the like.

"Optionally substituted heteroaralkyloxy means a radical -O-R<sup>a</sup> where R<sup>a</sup> is a heteroaralkyl radical e.g. 2-(pyridin-3-yl)ethoxy, 2-[3(2H)-pyridazon-1-yl]ethoxy and the like.

5

"Optional" or "optionally" means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "aryl group optionally mono- or di-

10 substituted with an alkyl group" means that the alkyl may but need not be present, and the description includes situations where the aryl group is mono- or disubstituted with an alkyl group and situations where the heterocyclo group is not substituted with the alkyl group.

15

"Amino protecting group" refers to those organic groups intended to protect nitrogen atoms against undesirable reactions during synthetic procedures e.g., benzyl, benzyloxycarbonyl (CBZ), *tert*-butoxycarbonyl (Boc), trifluoroacetyl, and the like.

The compounds of this invention may possess one or more

20 asymmetric centers; such compounds can therefore be produced as individual (*R*)- or (*S*)- stereoisomers or as mixtures thereof. Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for

25 the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see discussion in Chapter 4 of "Advanced Organic Chemistry", 4th edition J. March, John Wiley and Sons, New York, 1992).

30

A "pharmaceutically acceptable excipient" means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes an excipient that is acceptable for veterinary use as well as human pharmaceutical use. "A pharmaceutically acceptable excipient" as used in

- 12 -

the specification and claims includes both one and more than one such excipient.

A "pharmaceutically acceptable salt" of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include:

(1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynapthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or

(2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, tromethamine, *N*-methylglucamine, and the like.

- 13 -

"Pro-drugs" means any compound which releases an active parent drug according to Formula (I) *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of Formula (I) are prepared by modifying functional groups present in the compound of Formula (I) in such a way that the modifications may be cleaved *in vivo* to release the parent compound. Prodrugs include compounds of Formula (I) wherein a hydroxy, amino, or sulfhydryl group in compound (I) is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to esters (e.g., acetate, formate, and benzoate derivatives), carbamates (e.g., N,N-dimethylaminocarbonyl) of hydroxy functional groups in compounds of Formula (I), and the like.

"Treating" or "treatment" of a disease includes:

- (1) preventing the disease, i.e. causing the clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease,
- (2) inhibiting the disease, i.e., arresting or reducing the development of the disease or its clinical symptoms, or
- (3) relieving the disease, i.e., causing regression of the disease or its clinical symptoms.


A "therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

- 14 -

**NOMENCLATURE**



The naming and numbering of the compounds of this invention is illustrated below.

5





The nomenclature used in this application is generally based on the IUPAC recommendations, e.g., a compound of formula (I):

10

where  $R^1, R^2, R^4, R^6$  are hydrogen,  is 4-(3-hydroxypropyl)phenyl and  is 4-fluorophenyl is named 5-amino-1-(4-fluorophenyl)-4-[4-(3-hydroxypropyl)-benzoyl]pyrazole.

15


where  $R^1, R^2, R^4, R^6$  are hydrogen,  is 3-[3-(morpholin-4-yl)prop-1-ynyl]-phenyl and  is 4-fluorophenyl is named 5-amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylprop-1-ynyl)benzoyl]pyrazole.





- 15 -


Representative compounds of this invention are as follows:


- I. Compounds of Formula (I) where  $R^1$ ,  $R^2$  and  $R^4$  are hydrogen, B is phenyl and the other groups are as defined below are:
- 5 II.





CPD #	$R^3$ 	$R^5$	$R^6$	M. Pt. °C	Mass Spec. (M <sup>+</sup> )
1	4-(3-hydroxypropyl)phenyl	H	H	148-152.5	
2	4-[3-(morpholin-4-yl)prop-1-enyl]phenyl	4-F	H	166.7-168.2	
3	3-[3-(morpholin-4-yl)prop-1-ynyl]phenyl	4-F	H	145.2-145.8	
4	3-[3-(morpholin-4-yl)prop-1-enyl]phenyl	4-F	H	161.6-162.8	
5	4-[3-(morpholin-4-yl)propyl]phenyl	4-F, HCl	H		408
6	3-[3-(morpholin-4-yl)propyl]phenyl	4-F, HCl	H	211.9-212.6	
7	3-(3-hydroxyprop-1-ynyl)phenyl	4-F	H	219.2-219.6	
8	3-[3-(4-methylpiperazin-1-yl)prop-1-ynyl]phenyl	4-F, 2 HCl	H	270.8-271.1	
9	3-[3-(piperidin-1-yl)prop-1-ynyl]phenyl	4-F, HCl	H	205.3-207.4	
10	3-(2-aminosulfonylethyl)phenyl	4-F	H		370
11	3-cyanomethoxyphenyl	4-F	H		
12	3-[3-dimethylaminoprop-1-ynyl]phenyl	4-F, HCl	H	229.9-230	
13	3-[2-(morpholin-4-yl)ethoxy]phenyl	4-F	H		
14	3-[2-(morpholin-4-yl)ethoxy]phenyl	4-F, HCl	H	191.6-192.5	
15	3-[2-(morpholin-4-yl)ethoxy]phenyl	2-F	4-F		428
16	3-(3-methylaminoprop-1-ynyl)phenyl	4-F	H	210.6-210.8	
17	3-(4-methylpiperazin-1-yl)phenyl	2-F, HCl	H	142.5-151	
18	3-(morpholin-4-ylmethylcarbonyl)phenyl	4-F, HCl	H		408
19	3-(formamidoxime)phenyl	4-F	H	227-231.8	


CPD #	$R^3$ 	$R^5$	$R^6$	M. Pt. °C	Mass Spec. (M <sup>+</sup> )
20	3-(pyridin-3-yl)phenyl	4-F	H	222.4-223.0	
21	3-[3-(piperidin-1-yl)propoxy]phenyl	4-F	H		422
22	3-[2-(piperidin-1-yl)ethoxy]phenyl	4-F	H		408
23	3-[3-(morpholin-4-yl)propoxy]phenyl	4-F	H		424
24	3-[3-(4-methylpiperazin-1-yl)propoxy]phenyl	4-F	H		437
25	3-[3-(2-hydroxymethylpyrrolidin-1-yl)propoxy]-phenyl	4-F	H		438
26	3-[3-(2-aminocarbonylpyrrolidin-1-yl)propoxy]-phenyl	4-F	H		
27	3-(3-cyanopropoxy)phenyl	4-F	H		364
28	3-(3-diethylaminopropoxy)phenyl	4-F	H		410
29	2-[3-(4-methylpiperazin-1-yl)propoxy]phenyl	4-F	H		437
30	3-[3-(4-methylpiperazin-1-yl)propyl]phenyl	4-F . 2HCl	H	254.5-254.9	
31	3-[3-(4-methylpiperazin-1-yl)ethyl]phenyl	4-F . 2HCl	H	272.9-273.9	
32	3-[3-(piperidin-1-yl)propyl]phenyl	4-F . HCl	H	227.1-277.7	
33	3-(4-benzylpiperidin-1-yl)phenyl	4-F	H	156.2-160.9	
34	3-(methylaninocarbonylmethyloxy)phenyl	4-F	H	195.6-196.3	
35	3-(morpholin-4-ylcarbonylmethyloxy)phenyl	4-F	H	124.3-126.6	
36	3-(piperidin-1-ylcarbonylmethyloxy)phenyl	4-F	H		422
37	3-(diethylaminocarbonylmethyloxy)phenyl	4-F	H		410
38	3-(4-methylpiperazin-1-ylcarbonylmethyloxy)phenyl	4-F	H		437
39	3-(2-dimethylaminoethoxy)phenyl	4-F	H		368

CPD #	$R^3$ 	$R^5$	$R^6$	M. Pt. °C	Mass Spec. (M <sup>+</sup> )
40	3-(methylcarbonylmethoxy)phenyl	4-F	H		353
41	3-[4-(2-hydroxyethyl)piperazin-1-ylcarbonylmethoxy]-phenyl	4-F	H		467
42	3-[3-( <i>RS</i> )-hydroxypyrrolidin-1-ylcarbonylmethoxy]-phenyl	4-F	H		438
43	3-aminophenyl	4-F	H	165.2-165.8	
44	3-(3-imidazol-1-ylpropoxy)phenyl	4-F	H		405
45	3-[3-(4-( <i>RS</i> )-hydroxypiperidin-1-yl)propoxy]phenyl	4-F	H		438
46	3-(piperazin-1-yl)phenyl	4-F, 2 HCl	H	163.2-163.6	
47	3-(2-aminoethyl)phenyl	4-F, HCl	H	240.6-240.8	
48	3-[3-(morpholin-4-yl)propylamino]phenyl	4-F	H		423
49	3-[2-(morpholin-4-yl)ethylamino]phenyl	4-F	H		409
50	3-(2-aminosulfonylethyl)phenyl	4-F	H	170-170.4	
51	3-(piperidin-3-yl)phenyl	4-F	H	222.4-223	
52	3-(3-dimethylaminopropylamino)phenyl	4-F	H		381
53	3-[2-(3-( <i>RS</i> )-hydroxypyrrolidin-1-yl)ethylamino]phenyl	4-F	H		423
54	3-(2-hydroxyethoxy)phenyl	4-F	H		341
55	3-[3-(morpholin-4-yl)prop-1-ynyl]phenyl	2-F	4-F	190.4-191.2	
56	3-[3-(morpholin-4-yl)propyl]phenyl	2-F	4-F	226.4-227.5	
57	3-(2-aminopyridin-5-yl)phenyl	2-F, HCl	4-F	281.2-281.6	
58	3-(pyrimidin-3-yl)phenyl	4-F	H	237-241	

CPD #	$R^3$ 	$R^5$	$R^6$	M. Pt. °C	Mass Spec. (M <sup>+</sup> )
59	3-(1-methylpyridin-3-yl)phenyl.iodide	4-F	H	191.1-192.4	
60	3-(N-oxidopyridin-3-yl)phenyl	2-F	4-F	251.1-251.7	
61	3-(pyridin-4-yl)phenyl	2-F.HCl	4-F	218-226	376
62	3-(2-aminopyrimidin-5-yl)phenyl	2-F.HCl	4-F	272-275	
63	3-(2-amino-6-methylpyridin-5-yl)phenyl	2-F.HCl	4-F	215-217	
64	3-(6-methylpyridin-2-yl)phenyl	2-F.HCl	4-F	268-279	
65	3-(pyridin-3-yl)phenyl	4-Me.HCl	H	281.3-282.8	
66	3-(pyridin-3-yl)phenyl	H.HCl	3-MeO	256.1-256.5	
67	3-(pyridin-3-yl)phenyl	3-HO.HCl	H	269-273	
68	3-(pyridin-3-yl)phenyl	4-sulfamoyl	H.HCl	>300	419
69	3-(pyridin-3-yl)phenyl	2-Me.HCl	4-Me	250.7-251.8	
70	3-(N-oxidopyridin-3-yl)phenyl	2-Me	H	190.5-191.2	
71	3-(N-oxidopyridin-3-yl)phenyl	4-Me	H	212-213.5	
72	3-(2,6-dimethylpyridin-3-yl)phenyl	2-F.HCl	4-F	>300	440.879
73	3-(pyridin-3-yl)phenyl	2-Me.HCl	4-Cl		425.317
74	3-(pyridin-3-yl)phenyl	3-Me.HCl	4-Me		404.899
75	3-(pyridin-3-yl)phenyl	4-methylsulfonyl	H.HCl		454.936
76	3-(pyridin-3-yl)phenyl	2-Et.HCl	H		404.899
77	3-(imidazol-2-yl)phenyl	2-F.HCl	4-F		

CPD #		R <sup>5</sup>	R <sup>6</sup>	M. Pt. °C	Mass Spec. (M <sup>+</sup> )
78	3-(3-ethoxycarbonylphenyl)phenyl	4-F	H	267.4-268.7	
79	3-(pyridin-3-yl)phenyl	4-OH.HCl	H		
80	3-(3-carboxyphenyl)phenyl	4-F	H	117-128	429
81	3-{2-(piperidin-1-ylethoxy)}phenyl	4-F.HCl	H	210.2-211.2	
82	3-(pyridin-2-ylmethoxy)phenyl	4-F	H	176.1-177.3	
83	3-isopropylaminocarbonyloxyphenyl	4-F	H	225.2-230.1	
84	3-ethylaminocarbonyloxyphenyl	4-F	H	201.2-202.8	
85	3-(1,2-dihydroxyethyl)phenyl	4-F	H	52-56	341.34
86	3-(1-piperidinylmethyl)phenyl	2-F	4-F	116.6-118.6	396.439
87	3-(3-hydroxy-3-methyl-but-1-ynyl)phenyl	4-F	H	152.6-153.1	
88	3-(3-pyridylethynyl)phenyl	2-F	4-F	183.5-184.0	
89	3-{3-(S,S-dioxo-thiomorpholin-4-yl)-1-propynyl}phenyl	2-F	4-F	190.1-191.2	
90	3-(3-hydroxy-3-methylbutyl)phenyl	4-F	H	101.3-102.8	
91	3-(3-pyridylethyl)phenyl	2-F	4-F	153.4-153.7	
92	3-{3-(S,S-dioxo-thiomorpholin-4-yl)propyl}phenyl	2-F	4-F	123.8-125.9	
93	3-{2-(1-hydroxycyclopentyl)ethyl}phenyl	4-F	H	128.4-129.7	
94	3-{2-(1-hydroxycyclopentyl)ethynyl}phenyl	2-F	4-F	176.8-177.1	
95	3-{2-(1-hydroxycyclopentyl)ethyl}phenyl	2-F	4-F	120.3-121	
96	3-(3-hydroxybutyl)phenyl	2-F	4-F	111.1-112.6	
97	3-{2-(morpholin-4-yl)ethoxy}phenyl	2-F	H	130.8-135.1	

CPD #	<div> <div>R<sup>3</sup></div> <div>  </div> </div>	R <sup>5</sup>	R <sup>6</sup>	M. Pt. °C	Mass Spec. (M <sup>+</sup> )
98	3-[2-(morpholin-4-yl)ethoxy]phenyl	2-Cl	6-Cl	144.2-145.1	
99	3-(pyridin-3-yl)phenyl	2-F, HCl	4-F	256.5-257.7	
100	3-(2-methylsulfonyl)phenyl	4-F	H	151.5-155.6	
101	3-(2-methylsulfonyl)phenyl	2-F	4-F	157.2-157.7	
102	3-(2-ethylsulfonyl)phenyl	2-F	4-F	101.6-105.6	
103	3-(1,2-dihydroxyethyl)phenyl	2-F	4-F	59-64	
104	3-(2,2-dimethyl-1,3-dioxolan-5-yl)phenyl	4-F	H	94.5-100	
105	3-hydroxymethylphenyl	2-F	4-F	155.4-156.5	
106	3-[2(R),3-dihydroxypropoxy]phenyl	4-F	H	150.2-153.0	
107	3-[2(S),3-dihydroxypropoxy]phenyl	4-F	H	149.9-153.0	
108	3-(2-hydroxyethylsulfonyl)phenyl	2-F	4-F	92.1-93.8	
109	3-(1,2-dihydroxyethyl)phenyl	2-Me	H	83.0-85.5	
110	3-[2-(N-oxidomorpholin-4-yl)ethoxy]phenyl	4-F	H	185-186dec	
111	3-[2-(morpholin-4-yl)ethoxy]phenyl	2-F	6-F	178.9-181.2	
112	3-(2,3-dihydroxypropyl)phenyl	2-F	4-F	140-142	
113	3-(2-hydroxyethoxy)phenyl	4-F	H	165.3-166.7	
114	2-thienyl	4-F	H		287
115	2-furyl	4-F	H		271
116	2-methyl-3-furyl	2-F	4-F		
117	6-quinoliny	2-F	4-F	220-259.2	

CPD #		R <sup>5</sup>	R <sup>6</sup>	M. Pt. °C	Mass Spec. (M <sup>+</sup> )
118	2-hydroxyethylphenyl	4-F	H	154.5-155.0	
119	3-carboxymethoxyphenyl	4-F	H	215.9-216.2	
120	3-(pyridin-3-yl)phenyl	2-Me	H	223.7-225.1	
121	3-(3-sulfamoylphenyl)phenyl	4-F	H	214.6-217.5	
122	3-(3-methylsulfonylphenyl)phenyl	4-F	H		
123	3-methylsulfonylmethylphenyl	4-F	H		
124	3-sulfamoylmethylphenyl	4-F	H		
125	3-carboxymethylphenyl	4-F	H		
126	3-(2-hydroxyethylsulfonyl)phenyl	4-F	H		
127	3-[(2,2-dimethyl-1,3-dioxolan-5-yl)methoxy]phenyl	4-F	H		
128	3-oxiranylmethoxyphenyl	4-F	H		



While the broadest definition of this invention is set forth above, certain compounds of Formula (I) are preferred.

5 I.1. A preferred group of compounds of formula (I) is wherein  
R<sup>1</sup> is hydrogen or acyl;  
R<sup>2</sup> is hydrogen or alkyl;  
A is an aryl or heteroaryl ring  
and the other residues are as defined above.

10 I.2. Another preferred group of compounds of Formula (I) is  
wherein

R<sup>1</sup> is hydrogen, acyl or -P(O)(OH)<sub>2</sub>;  
R<sup>2</sup> is hydrogen, halo, alkyl or alkylthio;  
A is an aryl, heteroaryl or a heterocyclyl ring optionally fused  
15 to a phenyl ring provided that the heterocyclyl ring is  
attached to the carbonyl group via a carbon ring atom;  
B is an aryl or heteroaryl ring;  
R<sup>3</sup> is selected from the group consisting of:

- 20 (a) amino;
- (b) acylamino;
- (c) optionally substituted heterocycle;
- (d) heteroaryl optionally substituted with a  
substituent selected from halo, alkyl or alkoxy;
- (e) heteroalkyl;
- 25 (f) heteroalkenyl;
- (g) heteroalkynyl;
- (h) heteroalkoxy;
- (i) heteroalkylamino;
- (j) optionally substituted heterocyclalkyl;
- 30 (k) optionally substituted heterocyclalkenyl;
- (l) optionally substituted heterocyclalkynyl;
- (m) optionally substituted heterocyclalkoxy;
- (n) optionally substituted heterocyclalkylamino;
- (o) optionally substituted heterocyclalkylcarbonyl;

- 24 -

- (p) heteroalkylcarbonyl;
- (q)  $\text{-NHSO}_2\text{R}^6$  where  $\text{R}^6$  is alkyl, heteroalkyl or optionally substituted heterocyclalkyl;
- (r)  $\text{-NHSO}_2\text{NR}^7\text{R}^8$  where  $\text{R}^7$  and  $\text{R}^8$  are,  
 5 independently of each other, hydrogen, alkyl or heteroalkyl;
- (s)  $\text{-Y-(alkylene)-R}^9$  where:  
 Y is a single bond,  $\text{-O-}$ ,  $\text{-NH-}$  or  $\text{-S(O)}_n\text{-}$  (where n is an integer from 0 to 2); and  
 10  $\text{R}^9$  is cyano, heteroaryl,  $\text{-COOH}$ ,  $\text{-COR}^{10}$ ,  $\text{-COOR}^{11}$ ,  $\text{-CONR}^{12}\text{R}^{13}$ ,  $\text{-SO}_2\text{R}^{14}$ ,  $\text{-SO}_2\text{NR}^{15}\text{R}^{16}$ ,  $\text{-NHSO}_2\text{R}^{17}$  or  $\text{-NHSO}_2\text{NR}^{18}\text{R}^{19}$  where  $\text{R}^{10}$  is alkyl or optionally substituted heterocycle,  $\text{R}^{11}$  is alkyl, and  $\text{R}^{12}$ ,  $\text{R}^{13}$ ,  $\text{R}^{14}$ ,  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{17}$ ,  $\text{R}^{18}$  and  $\text{R}^{19}$  are, independently of  
 15 each other, hydrogen, alkyl or heteroalkyl;
- (t)  $\text{-C(=NR}^{20}\text{)(NR}^{21}\text{R}^{22})$  where  $\text{R}^{20}$ ,  $\text{R}^{21}$  and  $\text{R}^{22}$  independently represent hydrogen, alkyl or hydroxy, or  $\text{R}^{20}$  and  $\text{R}^{21}$  together are  $\text{-(CH}_2\text{)}_n\text{-}$  where n is 2 or 3 and  $\text{R}^{22}$  is hydrogen or alkyl;
- (u)  $\text{-NHC(X)NR}^{23}\text{R}^{24}$  where X is  $\text{-O-}$  or  $\text{-S-}$ , and  $\text{R}^{23}$  and  $\text{R}^{24}$  are, independently of each other,  
 20 hydrogen, alkyl or heteroalkyl;
- (v)  $\text{-CONR}^{25}\text{R}^{26}$  where  $\text{R}^{25}$  and  $\text{R}^{26}$  independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclalkyl, or  $\text{R}^{25}$  and  $\text{R}^{26}$  together with the nitrogen to which they are attached form an optionally substituted heterocycl ring;
- (w)  $\text{-S(O)}_n\text{R}^{27}$  where n is an integer from 0 to 2, and  
 25  $\text{R}^{27}$  is alkyl, heteroalkyl, optionally substituted heterocyclalkyl or  
 30  $\text{-NR}^{28}\text{R}^{29}$  where  $\text{R}^{28}$  and  $\text{R}^{29}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;

$\text{R}^4$  is selected from the group consisting of:

- 25 -

- (a) hydrogen;
- (b) halo;
- (c) alkyl; and
- (d) alkoxy;

5 R<sup>5</sup> is selected from the group consisting of :

- (a) hydrogen;
- (b) halo;
- (c) alkyl;
- (d) haloalkyl;
- 10 (e) thioalkyl;
- (f) hydroxy;
- (g) amino;
- (h) alkylamino;
- (i) dialkylamino;
- 15 (j) heteroalkyl;
- (k) optionally substituted heterocycle;
- (l) optionally substituted heterocyclalkyl; and
- (m) optionally substituted heterocyclalkoxy;

R<sup>6</sup> is selected from a group consisting of:

- 20 (a) hydrogen;
- (b) halo;
- (c) alkyl; and
- (d) alkoxy

II.1. Within these groups and preferred groups a more preferred

25 subgroup is wherein R<sup>3</sup> is selected from:

- (a) optionally substituted heterocycl;
- (b) aryl or heteroaryl both optionally substituted  
with a substituent selected from halo, alkyl,  
amino, alkoxy, carboxy, lower alkoxy carbonyl,  
30 SO<sub>2</sub>R' (where R' is alkyl) or SO<sub>2</sub>NHR'R'' (where R'  
and R'' are independently hydrogen or alkyl);
- (c) heteroalkyl;
- (d) heteroalkenyl;
- (e) heteroalkylamino;

- 26 -

- (f) heteroalkoxy;
- (g) optionally substituted heterocyclalkyl or heterocycloxy;
- (h) optionally substituted heterocyclalkenyl;
- (i) optionally substituted heterocyclalkynyl;
- (j) optionally substituted heterocyclalkoxy;
- (k) optionally substituted heterocyclalkylamino;
- (l) optionally substituted heterocyclalkylcarbonyl;
- (s) -Y-(alkylene)-R<sup>9</sup> where Y is a single bond, -O- or -NH- and R<sup>9</sup> is optionally substituted heteroaryl, -CONR<sup>12</sup>R<sup>13</sup>, SO<sub>2</sub>R<sup>14</sup>, -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup> -NHSO<sub>2</sub>R<sup>17</sup> or -NHSO<sub>2</sub>NR<sup>18</sup>R<sup>19</sup> where R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> are independently of each other hydrogen, alkyl or heteroalkyl;
- (x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;
- (m) arylaminoalkylene or heteroarylaminomethylene; or
- (n) Z-alkylene-NR<sup>30</sup>R<sup>31</sup> where Z is -NH-, -N(alkyl)- or -O-, and R<sup>30</sup> and R<sup>31</sup> are independently of each other, hydrogen, alkyl or heteroalkyl.

Within the above preferred group, more preferred groups of compounds are those wherein, A and B are aryl, preferably phenyl.

Within the above preferred and more preferred groups, an even more preferred group of compounds is that wherein:

R<sup>1</sup> is hydrogen;

R<sup>2</sup> is hydrogen or alkyl, preferably hydrogen or methyl, more preferably hydrogen.

R<sup>4</sup> is hydrogen, halo or alkyl, preferably hydrogen, chloro, fluoro or methyl, more preferably hydrogen.

- 27 -

II.1.1. Within the above preferred and more preferred groups, a particularly preferred group of compounds is that wherein  $R^3$  is at the 3-position and is optionally substituted heteroaryl, preferably pyridinyl, N-oxidopyridinyl or pyridonyl.

5

II.1.2. Another particularly preferred group of compounds is that wherein  $R^3$  is at the 3-position and is optionally substituted phenyl, preferably sulfamoylphenyl, alkylsulfamoylphenyl, carboxyphenyl, carboxamidophenyl, alkoxycarbonylphenyl, alkylaminocarbonylphenyl or dialkylaminocarbonylphenyl.

10

II.1.3. A third particularly preferred group of compounds is that wherein

$R^3$  is at the 3-position and is selected from:

15

- (a) heteroalkyl;
- (b) heteroalkoxy;
- (c) heteroalkylamino;
- (d) optionally substituted heterocyclalkyl;
- (e) optionally substituted heterocyclalkoxy;
- (f) optionally substituted heterocyclalkylamino;
- (g)  $-Y-(alkylene)-R^9$  where Y is a single bond,  $-O-$  or  $-NH-$  and  $R^9$  is optionally substituted heteroaryl,  $-CONR^{12}R^{13}$ ,  $SO_2R^{14}$ ,  $-SO_2NR^{15}R^{16}$ ,  $-NHSO_2R^{17}$  or  $-NHSO_2NR^{18}R^{19}$  where  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  are independently of each other hydrogen, alkyl or heteroalkyl; or
- (h)  $Z-alkylene-NR^{30}R^{31}$  where Z is  $-NH-$ ,  $-N(alkyl)-$  or  $-O-$ , and  $R^{30}$  and  $R^{31}$  are independently of each other, hydrogen, alkyl or heteroalkyl.

20

25

30

Preferred groups for  $R^3$  include amino, 3-dimethylaminopropoxy, 2-dimethylaminoethoxy, 2-dimethylaminoethyl, 3-dimethylaminopropyl, 4-dimethylaminobutyl, 2-dimethylaminoethylamino, 3-dimethylaminopropylamino, 3-dimethylaminoprop-1-enyl, 3-

- 28 -

dimethylaminoprop-1-ynyl and 2-dimethylaminoethylcarbonyl, preferably amino.

Another group of preferred groups for  $R^3$  is selected from 3-(morpholin-4-yl)propoxy, 2-(morpholin-4-yl)ethoxy, 3-(morpholin-4-yl)propyl, 2-(morpholin-4-yl)ethyl, 4-(morpholin-4-yl)butyl, 3-(morpholin-4-yl)propylamino, 2-(morpholin-4-yl)-ethylamino, 3-(morpholin-4-yl)-prop-1-enyl, 3-(morpholin-4-yl)prop-1-ynyl, 4-methylpiperazin-1-yl, piperazin-1-yl, pyridin-3-yl, morpholin-4-ylmethylcarbonyl, 3-dimethylaminoprop-1-enyl, 3-dimethylaminoprop-1-ynyl, 2-aminosulfonylethyl, 2-aminosulfonylethenyl, acetylamino and trifluoroacetylamino, preferably 2-(morpholin-4-yl)ethoxy and 3-(morpholin-4-yl)-propyl.

A fourth group of particularly preferred compounds is that where  $R^5$  is halo or alkyl and  $R^6$  is hydrogen, halo or alkyl, preferably  $R^5$  is 4-F or 2-Me and  $R^6$  is hydrogen, or  $R^5$  is 2-F and  $R^6$  is 4-F.

III. Within the above preferred and more preferred I and II groups, an even more preferred group of compounds is that wherein  $R^1$  and  $R^2$  are hydrogen and B is phenyl; particularly wherein A is phenyl; more particularly wherein  $R^4$  is hydrogen; and  $R^5$  is halo or alkyl, particularly wherein  $R^5$  is chloro, fluoro or methyl and  $R^6$  is hydrogen, chloro, fluoro, methyl or methoxy.

IV.1. Within the above preferred and more preferred  $R^3$  groups in I, II and III, a particularly preferred  $R^3$  group is that wherein  $R^3$  is optionally substituted heteroaryl, particularly wherein  $R^3$  is pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, N-oxidopyridin-2-yl, N-oxidopyridin-3-yl, N-oxidopyridin-4-yl or pyridon-2-yl, all optionally substituted, especially wherein  $R^3$  is at the 3-position, particularly together with  $R^5$  and  $R^6$  wherein  $R^5$  is 4-F or 2-Me and  $R^6$  is hydrogen;

IV.2. Within the above preferred and even more preferred  $R^3$  groups I, II and III, another particular preferred  $R^3$  group is that wherein  $R^3$  is

- 29 -

optionally substituted phenyl, particularly wherein  $R^3$  is 3-sulfamoylphenyl, 3-methylsulfonylphenyl, 3-carboxyphenyl or 3-ethoxycarbonylphenyl, particularly wherein  $R^3$  is at the 3-position, especially in combination wherein  $R^5$  is 4-F or 2-Me and  $R^6$  is hydrogen.

5

IV.3. Within the above preferred and even more preferred  $R^3$  groups in I, II and III, another particular preferred group of  $R^3$  groups are those wherein  $R^3$  is:

- (a) heteroalkyl;
- 10 (b) heteroalkoxy;
- (c) heteroalkylamino;
- (d) optionally substituted heterocyclalkyl;
- (e) optionally substituted heterocyclalkoxy;
- (f) optionally substituted heterocyclalkylamino;
- 15 (g)  $-Y-(alkylene)-R^9$  where Y is a single bond, -O- or -NH- and  $R^9$  is optionally substituted heteroaryl,  $-\text{CONR}^{12}\text{R}^{13}$ ,  $\text{SO}_2\text{R}^{14}$ ,  $-\text{SO}_2\text{NR}^{15}\text{R}^{16}$ ,  $-\text{NHSO}_2\text{R}^{17}$  or  $-\text{NHSO}_2\text{NR}^{18}\text{R}^{19}$  where  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  are independently of each other hydrogen, alkyl or heteroalkyl; or
- 20 (h)  $Z-(alkylene)-\text{NR}^{30}\text{R}^{31}$  where Z is -NH-, -N(alkyl)- or -O-, and  $R^{30}$  and  $R^{31}$  are independently of each other, hydrogen, alkyl or heteroalkyl.

- a) A more preferred subgroup within these particular preferred  $R^3$  groups is that wherein  $R^3$  is heteroalkyl, especially wherein  $R^3$  is at the 3-position and is selected from the group consisting of 2-dimethylaminoethyl, 3-dimethylaminopropyl, 4-dimethylaminobutyl, 2-dimethylaminoethylamino, 3-dimethylaminopropylamino, hydroxymethyl, 1,2-dihydroxyethyl, 3-hydroxy-3-methyl-1-butyl or 3-hydroxybutyl, particularly in combination with  $R^5$  and  $R^6$ , wherein  $R^5$  is 2-F and  $R^6$  is 4-F, or wherein  $R^5$  is 4-F and  $R^6$  is hydrogen or wherein  $R^5$  is 2-Me and  $R^6$  is hydrogen.

30

- 30 -

b) Another more preferred subgroup within these particular preferred  $R^3$  groups is that wherein  $R^3$  is heteroalkoxy or heteroalkylamino, particularly wherein  $R^3$  is at the 3-position and is selected from the group consisting of 3-dimethylaminopropoxy, 2-dimethylaminoethoxy, 2-hydroxyethoxy, 2,3-dihydroxypropoxy, 2-dimethylaminoethylamino and 3-dimethylaminopropylamino, especially in combination with  $R^5$  and  $R^6$  wherein  $R^5$  is 4-F or 2-Me and  $R^6$  is hydrogen.

c) Another preferred  $R^3$  subgroup within these particular preferred groups is that wherein  $R^3$  is optionally substituted heterocyclalkyl, optionally substituted heterocyclalkoxy or optionally substituted heterocyclalkylamino, particularly wherein  $R^3$  is at the 3-position and is selected from the group consisting of 3-(morpholin-4-yl)propoxy, 2-(morpholin-4-yl)ethoxy, 2-(2-oxo-pyrrolidin-1-yl)ethoxy, 3-(morpholin-4-yl)propyl, 2-(morpholin-4-yl)ethyl, 4-(morpholin-4-yl)butyl, 3-(morpholin-4-yl)propylamino, 2-(morpholin-4-yl)ethylamino, 4-hydroxypiperidinylmethyl, 2-(S,S-dioxo-thiamorpholin-4-yl)ethyl, 3-(S,S-dioxo-thiamorpholin-4-yl)propyl and N-methylpiperazinylmethyl, especially in combination with  $R^5$  and  $R^6$  wherein  $R^5$  is 4-F or 2-Me and  $R^6$  is hydrogen.

d) Another preferred  $R^3$  subgroup is that wherein  $R^3$  is -Y-(alkylene)- $R^9$  where Y is a single bond, -O- or -NH- and  $R^9$  is optionally substituted heteroaryl, -CONR<sup>12</sup>R<sup>13</sup>, SO<sub>2</sub>R<sup>14</sup>, -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup> -NHSO<sub>2</sub>R<sup>17</sup> or -NHSO<sub>2</sub>NR<sup>18</sup>R<sup>19</sup> where R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> are independently of each other hydrogen, alkyl or heteroalkyl, particularly wherein Y is a single bond and  $R^9$  is SO<sub>2</sub>R<sup>14</sup> or -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup>. Preferably  $R^3$  is methylsulfonylethyl or sulfamoylethyl and  $R^3$  is preferably combined with  $R^5$  and  $R^6$  wherein  $R^5$  is 4-F or 2-Me and  $R^6$  is hydrogen.

V. It has to be noted that in the embodiment I.2. of compounds of formula (I) wherein

$R^1$  is hydrogen, acyl or -P(O)(OH)<sub>2</sub>;

$R^2$  is hydrogen, halo, alkyl or alkylthio;



- 31 -

A is an aryl, heteroaryl or a heterocyclyl ring optionally fused to a phenyl ring provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom;

B is an aryl or heteroaryl ring;

R<sup>3</sup> is selected from the group consisting of:

- (a) amino;
- (b) acylamino;
- (c) optionally substituted heterocycle;
- (d) heteroaryl optionally substituted with a  
substituent selected from halo, alkyl or alkoxy;
- (e) heteroalkyl;
- (f) heteroalkenyl;
- (g) heteroalkynyl;
- (h) heteroalkoxy;
- (i) heteroalkylamino;
- (j) optionally substituted heterocyclylalkyl;
- (k) optionally substituted heterocyclylalkenyl;
- (l) optionally substituted heterocyclylalkynyl;
- (m) optionally substituted heterocyclylalkoxy;
- (n) optionally substituted heterocyclylalkylamino;
- (o) optionally substituted heterocyclylalkylcarbonyl;
- (p) heteroalkylcarbonyl;
- (q) -NHSO<sub>2</sub>R<sup>6</sup> where R<sup>6</sup> is alkyl, heteroalkyl or  
optionally substituted heterocyclylalkyl;
- (r) -NHSO<sub>2</sub>NR<sup>7</sup>R<sup>8</sup> where R<sup>7</sup> and R<sup>8</sup> are,  
independently of each other, hydrogen, alkyl or  
heteroalkyl;
- (s) -Y-(alkylene)-R<sup>9</sup> where:  
Y is a single bond, -O-, -NH- or -S(O)<sub>n</sub>- (where n is  
an integer from 0 to 2); and  
R<sup>9</sup> is cyano, heteroaryl, -COOH, -COR<sup>10</sup>, -COOR<sup>11</sup>,  
-CONR<sup>12</sup>R<sup>13</sup>, -SO<sub>2</sub>R<sup>14</sup>, -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup>, -NHSO<sub>2</sub>R<sup>17</sup> or  
-NHSO<sub>2</sub>NR<sup>18</sup>R<sup>19</sup> where R<sup>10</sup> is alkyl or optionally  
substituted heterocycle, R<sup>11</sup> is alkyl, and R<sup>12</sup>, R<sup>13</sup>,

- 32 -

$R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$  and  $R^{19}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;

- (t)  $-C(=NR^{20})(NR^{21}R^{22})$  where  $R^{20}$ ,  $R^{21}$  and  $R^{22}$  independently represent hydrogen, alkyl or hydroxy, or  $R^{20}$  and  $R^{21}$  together are  $-(CH_2)_n-$  where  $n$  is 2 or 3 and  $R^{22}$  is hydrogen or alkyl;
- (u)  $-NHC(X)NR^{23}R^{24}$  where  $X$  is  $-O-$  or  $-S-$ , and  $R^{23}$  and  $R^{24}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;
- (v)  $-CONR^{25}R^{26}$  where  $R^{25}$  and  $R^{26}$  independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclalkyl, or  $R^{25}$  and  $R^{26}$  together with the nitrogen to which they are attached form an optionally substituted heterocycl ring;
- (w)  $-S(O)_nR^{27}$  where  $n$  is an integer from 0 to 2, and  $R^{27}$  is alkyl, heteroalkyl, optionally substituted heterocyclalkyl or  $-NR^{28}R^{29}$  where  $R^{28}$  and  $R^{29}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;

$R^4$  is selected from the group consisting of:

- (a) hydrogen;
- (b) halo;
- (c) alkyl; and
- (d) alkoxy;

$R^5$  is selected from the group consisting of :

- (a) hydrogen;
- (b) halo;
- (c) alkyl;
- (d) haloalkyl;
- (e) thioalkyl;
- (f) hydroxy;
- (g) amino;
- (h) alkylamino;

- 33 -

- (i) dialkylamino;
- (j) heteroalkyl;
- (k) optionally substituted heterocycle;
- (l) optionally substituted heterocyclalkyl; and
- (m) optionally substituted heterocyclalkoxy; and

$R^6$  is selected from a group consisting of:

- (a) hydrogen;
- (b) halo;
- (c) alkyl; and
- (d) alkoxy;

further preferred embodiments exist.

V.1. In one preferred embodiment a preferred subgroup is that wherein  $R^3$  is selected from:

- (a) optionally substituted heterocycle;
- (b) heteroaryl optionally substituted with a substituent selected from halo, alkyl or alkoxy;
- (c) optionally substituted heterocyclalkyl;
- (d) optionally substituted heterocyclalkenyl;
- (e) optionally substituted heterocyclalkynyl;
- (f) optionally substituted heterocyclalkoxy;
- (g) optionally substituted heterocyclalkylamino;
- (h) optionally substituted heterocyclalkylcarbonyl;
- (i) heteroalkenyl;
- (j) heteroalkynyl;
- (k)  $-Y-(alkylene)-R^9$  where Y is a single bond,  $-O-$  or  $-NH-$  and  $R^9$  is  $-CONR^{12}R^{13}$  or  $-SO_2NR^{15}R^{16}$  where  $R^{12}$ ,  $R^{13}$ ,  $R^{15}$  and  $R^{16}$  are independently of each other hydrogen, alkyl or heteroalkyl; and
- (l) acylamino,

particularly wherein  $R^2$  is hydrogen or alkyl and B is aryl;  
 particularly wherein  $R^4$  is hydrogen, halo or alkyl,  $R^5$  is hydrogen, halo, alkyl, hydroxy or heteroalkyl and  $R^6$  is hydrogen, halo or alkyl; particularly

- 34 -

wherein A is aryl; R<sup>2</sup> is hydrogen and B is phenyl; particularly wherein A is phenyl; particularly wherein R<sup>1</sup> is hydrogen, R<sup>4</sup> is hydrogen, chloro, fluoro or methyl, R<sup>5</sup> is hydrogen, fluoro, methyl, hydroxy, amino, hydroxymethyl or aminomethyl and R<sup>6</sup> is hydrogen or fluoro;

5 particularly wherein R<sup>3</sup> is at the 3-position and is selected from 3-(morpholin-4-yl)propoxy, 2-(morpholin-4-yl)ethoxy, 3-(morpholin-4-yl)propyl, 2-(morpholin-4-yl)ethyl, 4-(morpholin-4-yl)butyl, 3-(morpholin-4-yl)propylamino, 2-(morpholin-4-yl)ethylamino, 3-(morpholin-4-yl)prop-1-enyl, 3-(morpholin-4-yl)prop-1-ynyl, 4-methylpiperazin-1-yl, piperazin-1-yl,  
10 pyridin-3-yl, morpholin-4-ylmethyl-carbonyl, 3-dimethylaminoprop-1-enyl, 3-dimethylaminoprop-1-ynyl, 2-amino-sulfonylethyl, 2-aminosulfonylethenyl, acetylamino and trifluoroacetylamino.

V.2. Another preferred subgroup of R<sup>3</sup> is that wherein R<sup>3</sup> is selected from:

- 15 (a) amino;  
(b) heteroalkyl;  
(c) heteroalkoxy;  
(d) heteroalkylamino; and  
(e) heteroalkylcarbonyl;

20 particularly wherein R<sup>2</sup> is hydrogen or alkyl and B is aryl;  
particularly wherein R<sup>4</sup> is hydrogen, halo or alkyl, R<sup>5</sup> is hydrogen, halo, alkyl, hydroxy or heteroalkyl and R<sup>6</sup> is hydrogen, halo or alkyl; particularly wherein A is aryl, R<sup>2</sup> is hydrogen and B is phenyl; particularly wherein A is phenyl; particularly wherein R<sup>1</sup> is hydrogen, R<sup>4</sup> is hydrogen, chloro, fluoro  
25 or methyl, R<sup>5</sup> is hydrogen, fluoro, methyl, hydroxy, amino, hydroxymethyl or aminomethyl and R<sup>6</sup> is hydrogen or fluoro;

particularly wherein R<sup>3</sup> is at the 3-position and is selected from amino, 3-dimethylaminopropoxy, 2-dimethylaminoethoxy, 2-dimethylaminoethyl, 3-dimethylaminopropyl, 4-dimethylaminobutyl, 2-dimethylaminoethylamino, 3-dimethylaminopropylamino and 2-dimethylaminoethylcarbonyl.  
30

V.3. In another preferred group of compounds of formula (I) is that

- 35 -

wherein A is a phenyl ring fused to a 5 or 6 membered monocyclic saturated ring optionally containing 1 or 2 heteroatoms independently selected from nitrogen or oxygen the remaining ring atoms being carbon, where one or two carbon atoms are optionally replaced by a carbonyl group; particularly wherein R<sup>2</sup> is hydrogen or alkyl and B is aryl; particularly  
5 wherein R<sup>4</sup> is hydrogen, halo or alkyl, R<sup>5</sup> is hydrogen, halo, alkyl, hydroxy or heteroalkyl and R<sup>6</sup> is hydrogen, halo or alkyl; particularly wherein R<sup>2</sup> is hydrogen and B is phenyl;

particularly wherein R<sup>3</sup> is selected from:

- 10 (a) optionally substituted heterocycle;
- (b) heteroaryl optionally substituted with a substituent selected from halo, alkyl or alkoxy;
- (c) optionally substituted heterocyclalkyl;
- (d) optionally substituted heterocyclalkenyl;
- 15 (e) optionally substituted heterocyclalkynyl.
- (f) optionally substituted heterocyclalkoxy;
- (g) optionally substituted heterocyclalkylamino;
- (h) optionally substituted heterocyclalkylcarbonyl;
- (i) heteroalkenyl;
- 20 (j) heteroalkynyl;
- (k) -Y-(alkylene)-R<sup>9</sup> where Y is a single bond, -O- or -NH- and R<sup>9</sup> is -CONR<sup>12</sup>R<sup>13</sup> or -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup> where R<sup>12</sup>, R<sup>13</sup>, R<sup>15</sup> and R<sup>16</sup> are independently of each other hydrogen, alkyl or heteroalkyl; and
- 25 (l) acylamino or

wherein R<sup>3</sup> is selected from:

- (a) amino;
- (b) heteroalkyl;
- (c) heteroalkoxy;
- 30 (d) heteroalkylamino; and
- (f) heteroalkylcarbonyl;

more particularly wherein A is 2,3-dihydrobenzo[1,4]dioxane, chroman, isochroman, 2,3-dihydrobenzofuran, 1,3-dihydroisobenzofuran, benzo[1,3]dioxole, 1,2,3,4-tetrahydroisoquinoline, 1,2,3,4-

- 36 -

tetrahydroquinoline, 2,3-dihydro-1H-indole, 2,3-dihydro-1H-isoindole, benzimidazol-2-one or 3H-benzoxazolin-2-one.

V.4. Within the embodiment of the compounds of formula (I) wherein  
 5 R<sup>3</sup> is selected from:

- (a) optionally substituted heterocycle;
- (b) heteroaryl optionally substituted with a  
 substituent selected from halo, alkyl or alkoxy;
- (c) optionally substituted heterocyclalkyl;
- 10 (d) optionally substituted heterocyclalkenyl;
- (e) optionally substituted heterocyclalkynyl;
- (f) optionally substituted heterocyclalkoxy;
- (g) optionally substituted heterocyclalkylamino;
- (h) optionally substituted heterocyclalkylcarbonyl;
- 15 (i) heteroalkenyl;
- (j) heteroalkynyl;
- (k) -Y-(alkylene)-R<sup>9</sup> where Y is a single bond, -O- or -  
 NH- and R<sup>9</sup> is -CONR<sup>12</sup>R<sup>13</sup> or -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup> where  
 R<sup>12</sup>, R<sup>13</sup>, R<sup>15</sup> and R<sup>16</sup> are independently of each  
 20 other hydrogen, alkyl or heteroalkyl; and
- (l) acylamino;

and wherein R<sup>2</sup> is hydrogen or alkyl and B is aryl, and  
 wherein R<sup>4</sup> is hydrogen, halo or alkyl, R<sup>5</sup> is hydrogen, halo, alkyl, hydroxy  
 or heteroalkyl and R<sup>6</sup> is hydrogen, halo or alkyl, another preferred group is  
 25 that

wherein A is heteroaryl, R<sup>2</sup> is hydrogen and B is phenyl; particularly  
 wherein A is furanyl, thienyl or pyridyl; particularly  
 wherein R<sup>1</sup> is hydrogen, R<sup>4</sup> is hydrogen, chloro, fluoro or methyl, R<sup>5</sup> is  
 hydrogen, fluoro, methyl, hydroxy, amino, hydroxymethyl or aminomethyl  
 30 and R<sup>6</sup> is hydrogen or fluoro; particularly

wherein R<sup>3</sup> is selected from 3-(morpholin-4-yl)propoxy, 2-(morpholin-4-  
 yl)ethoxy, 3-(morpholin-4-yl)propyl, 2-(morpholin-4-yl)ethyl, 4-(morpholin-  
 4-yl)butyl, 3-(morpholin-4-yl)propylamino, 2-(morpholin-4-yl)ethylamino, 3-  
 (morpholin-4-yl)prop-1-enyl, 3-(morpholin-4-yl)prop-1-ynyl, 4-

methylnpiperazin-1-yl, piperazin-1-yl, pyridin-3-yl, morpholin-4-ylmethylcarbonyl, 3-dimethylaminoprop-1-enyl, 3-dimethylaminoprop-1-ynyl, 2-aminosulfonylethyl, 2-aminosulfonylethenyl, acetyl amino and trifluoroacetyl amino.

5

V.5. Another group of compounds of formula (I) wherein R<sup>3</sup> is selected from:

- (a) amino;
- (b) heteroalkyl;
- 10 (c) heteroalkoxy;
- (d) heteroalkylamino; and
- (e) heteroalkylcarbonyl, and

wherein R<sup>2</sup> is hydrogen or alkyl and B is aryl, and wherein R<sup>4</sup> is hydrogen, halo or alkyl, R<sup>5</sup> is hydrogen, halo, alkyl, hydroxy or heteroalkyl  
15 and R<sup>6</sup> is hydrogen, halo or alkyl, also such are preferred wherein A is heteroaryl, R<sup>2</sup> is hydrogen and B is phenyl; particularly wherein A is furanyl, thienyl or pyridyl; particularly

wherein R<sup>1</sup> is hydrogen, R<sup>4</sup> is hydrogen, chloro, fluoro or methyl, R<sup>5</sup> is hydrogen, fluoro, methyl, hydroxy, amino, hydroxymethyl or aminomethyl  
20 and R<sup>6</sup> is hydrogen or fluoro; particularly wherein R<sup>3</sup> is selected from amino, 3-dimethylamino-propoxy, 2-dimethylaminoethoxy, 2-dimethylaminoethyl, 3-dimethylaminopropyl, 4-dimethylaminobutyl, 2-dimethylaminoethylamino, 3-dimethylaminopropylamino and 2-dimethylaminoethylcarbonyl.

25

V.6. Another group of compounds of formula (I), wherein R<sup>3</sup> is selected from:

- (a) optionally substituted heterocycle;
- (b) heteroaryl optionally substituted with a  
30 substituent selected from halo, alkyl or alkoxy;
- (c) optionally substituted heterocyclalkyl;
- (d) optionally substituted heterocyclalkenyl;
- (e) optionally substituted heterocyclalkynyl;
- (f) optionally substituted heterocyclalkoxy;

- 38 -

- (g) optionally substituted heterocyclalkylamino;  
 (h) optionally substituted heterocyclalkylcarbonyl;  
 (i) heteroalkenyl;  
 (j) heteroalkynyl;  
 5 (k) -Y-(alkylene)-R<sup>9</sup> where Y is a single bond, -O- or -NH- and R<sup>9</sup> is -CONR<sup>12</sup>R<sup>13</sup> or -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup> where R<sup>12</sup>, R<sup>13</sup>, R<sup>15</sup> and R<sup>16</sup> are independently of each other hydrogen, alkyl or heteroalkyl; and  
 (l) acylamino; and
- 10 R<sup>2</sup> is hydrogen or alkyl and B is aryl and R<sup>4</sup> is hydrogen, halo or alkyl, R<sup>5</sup> is hydrogen, halo, alkyl, hydroxy or heteroalkyl and R<sup>6</sup> is hydrogen, halo or alkyl; is that,  
 wherein A is heterocycl ring, R<sup>2</sup> is hydrogen and B is phenyl;  
 particularly  
 15 wherein A is tetrahydrofuran, tetrahydropyran, dioxane or dioxolane, particularly, wherein R<sup>1</sup> is hydrogen, R<sup>4</sup> is hydrogen, chloro, fluoro or methyl, R<sup>5</sup> is hydrogen, fluoro, methyl, hydroxy, amino, hydroxymethyl or aminomethyl and R<sup>6</sup> is hydrogen or fluoro;  
 particularly, wherein R<sup>3</sup> is selected from amino, 3-dimethyl-  
 20 aminopropoxy, 2-dimethylaminoethoxy, 2-dimethylaminoethyl, 3-dimethylamino-propyl, 4-dimethylaminobutyl, 2-dimethylaminoethylamino, 3-dimethylaminopropyl-amino and 2-dimethylaminoethylcarbonyl.

25 Preferred compounds of the invention are selected from the group consisting of:

5-amino-1-(4-fluorophenyl)-4-[3-(2-morpholin-4-ylethoxy)benzoyl]pyrazole,  
 5-amino-1-(2,4-difluorophenyl)-4-[3-(3-morpholin-4-ylpropyl)benzoyl]-  
 30 pyrazole,  
 5-amino-4-(3-aminobenzoyl)-1-(4-fluorophenyl)pyrazole,  
 5-amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylpropyl)benzoyl]pyrazole,  
 5-amino-4-[3-(2-aminosulfonylethenyl)benzoyl]-1-(4-fluorophenyl)pyrazole,  
 5-amino-4-(3-acetylaminobenzoyl)-1-phenylpyrazole,



- 39 -


- 5-amino-4-[3-(2-aminoethyl)benzoyl]-1-(4-fluorophenyl)pyrazole,  
 5-amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylpropylamino)benzoyl]-  
 pyrazole,  
 5-amino-4-[3-(2-aminosulfonyl)ethyl]benzoyl]-1-(4-fluorophenyl)pyrazole,  
 5 5-amino-1-(4-fluorophenyl)-4-(3-pyridin-3-ylbenzoyl)pyrazole, and

Other preferred compounds of the invention are selected from the group consisting of:

- 10 5-amino-1-(2-methylphenyl)-4-[3-pyridin-3-yl]benzoyl]pyrazole,  
 5-amino-1-(2-methylphenyl)-4-[3-(N-oxidopyridin-3-yl)benzoyl]pyrazole,  
 5-amino-4-[3-(2,3-dihydroxypropoxy)benzoyl]-1-(4-fluorophenyl)pyrazole,  
 5-amino-4-[3-(1,2-dihydroxyethyl)benzoyl]-1-(4-fluorophenyl)pyrazole,  
 5-amino-1-(4-fluorophenyl)-4-[3-(sulfamoylbenzoyl]pyrazole,  
 15 5-amino-1-(4-fluorophenyl)-4-[3-(3-hydroxy-3-methylbutyl)benzoyl]-  
 pyrazole,  
 5-amino-1-(4-fluorophenyl)-4-[3-(2-(1-hydroxycyclopentyl)ethyl)benzoyl]-  
 pyrazole,  
 5-amino-4-[3-(2-methylsulfonyl)ethyl]benzoyl]-1-(4-fluorophenyl)pyrazole,  
 20 and  
 5-amino-1-(2,4-difluorophenyl)-4-[3-(2-hydroxyethylsulfonyl)benzoyl]-  
 pyrazole.


In a second aspect the present invention relates to processes for  
 25 preparing compounds of formula (I). In one embodiment the compounds of  
 formula (I) can be prepared by a process comprising

- (i) reacting a 2-keto-3-phenylaminoacrylonitrile of Formula 1:




- 40 -

with a hydrazine of Formula 2:




5        where R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are as defined above to provide a compound of Formula (I) where R<sup>1</sup> is hydrogen; or

(ii)    reacting a 2-keto-3-phenylaminoacrylonitrile of formula 3:



10        where Z is either hydroxy, nitro or halo group and R<sup>4</sup> is as defined above with a hydrazine of formula 2 to provide a compound of formula 4:




15        followed by conversion of the Z group to the desired R<sup>3</sup> group to provide a compound of Formula (I) where R<sup>1</sup> is hydrogen;

- (iii)    optionally modifying any of the R<sup>1</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> or R<sup>6</sup> groups;
- (iv)    optionally converting the compound of Formula (I) prepared in Steps (i), (ii) or (iii) above, to the corresponding acid addition salt by treatment with an acid;
- 20        (v)    optionally converting the compound of Formula (I) prepared in Steps (i), (ii) or (iii) above, to the corresponding free base by treatment with a base; and

- 41 -

- (vi) optionally separating a mixture of stereoisomers of a compound of Formula (I) prepared in Steps (i) - (v) above, to give a single stereoisomer.

5 In another embodiment the compounds can be prepared by a process comprising reacting a compound of Formula 5:



10 where  $R^5$  and  $R^6$  are as defined above and L is a leaving group under organometallic displacement reaction conditions



with an organometallic reagent of formula  $R^3$   $R^4$   $M^+$  where  $R^3$  and  $R^4$  are as defined above and M is a metallic moiety to provide a compound of Formula (I) where  $R^1$  is hydrogen;

- 15 (ii) optionally modifying any of the  $R^1$ ,  $R^3$ ,  $R^4$ ,  $R^5$  or  $R^6$  groups;  
 (iii) optionally converting the compound of Formula (I) prepared in Steps (i) or (ii) above, to the corresponding acid addition salt by treatment with an acid;  
 (iv) optionally converting the compound of Formula (I) prepared in Steps  
 20 (i) or (ii) above, to the corresponding free base by treatment with a base; and  
 (v) optionally separating a mixture of stereoisomers of a compound of Formula (I) prepared in Steps (i) or (iv) above, to give a single stereoisomer.

25

More specifically, the compounds of this invention can be made by the methods depicted in the reaction schemes shown below.

- 42 -

The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, Wisconsin, USA), Bachem (Torrance, California, USA), Emka-Chemie, or Sigma (St. Louis, Missouri, USA) or are prepared by methods known to those skilled in the art following procedures set forth in references such as *Fieser and Fieser's Reagents for Organic Synthesis*, Volumes 1-17 (John Wiley and Sons, 1991); *Rodd's Chemistry of Carbon Compounds*, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989), *Organic Reactions*, Volumes 1-40 (John Wiley and Sons, 1991), *March's Advanced Organic Chemistry*, (John Wiley and Sons, 4th Edition), and *Larock's Comprehensive Organic Transformations* (VCH Publishers Inc., 1989). These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure.

The starting materials and the intermediates of the reaction may be isolated and purified if desired using conventional techniques, including but not limited to filtration, distillation, crystallization, chromatography, and the like. Such materials may be characterized using conventional means, including physical constants and spectral data.

#### Preparation of compounds of Formula (I)


Schemes A, B and C describe methods to generate the compounds of Formula (I).

#### Scheme A


Compounds of Formula (I) where R<sup>2</sup> is hydrogen and other groups are as defined in the Summary of the Invention are prepared as described below.

#### Method (a)

- 43 -

Method (b)

5



In general, compounds of Formula (I) can be prepared by following either method (a) or (b) as described below.

10

Method (a)

Reaction of a 2-ketoacetonitrile of formula 1 [where Z is halo (e.g., bromo or iodo), alkoxy, nitro or acetyl amino] with N,N-diphenylformamidine

- 44 -

gives a 2-keto-3-phenylamino-acrylonitrile of formula 2. The reaction occurs upon heating in a high boiling aromatic hydrocarbon such as toluene, xylene, and the like.

5        In general, compounds of formula 1 are either commercially available or they can be prepared by methods well known in the art. For example, 2-arylacetonitriles of formula 1 such as 4-methoxybenzoylacetonitrile, 3-nitrobenzoylacetonitrile are commercially available. Others can be prepared by treating acetonitrile with a base such as *n*-butyllithium followed by  
10       reaction of the formed acetonitrile anion with an aroyl/heteroaroyl halide or an aryl/heteroaryl ester as described in Sjogren, E. B., et al., *J. Med. Chem.*, 34, 3295, (1991).


15       Reaction of the 2-keto-3-phenylaminoacrylonitrile of formula 2 with a hydrazine of formula 3 provides a 5-amino-4-ketopyrazole of formula 4. This reaction is generally carried out in a polar solvent such as ethanol, isopropanol, and the like. Aryl/heteroaryl hydrazines of formula 2 such as 2- or 3-chlorophenylhydrazine, 2-,3-, or 4-fluorophenylhydrazine, phenylhydrazine, 2-hydrazinopyridine, 2-hydrazinobenzothiazole, 2-  
20       hydrazinoquinoline etc., are commercially available.

Compound 4 is then converted to a compound of Formula (I) where R<sup>1</sup> is hydrogen and R<sup>3</sup> is as defined in the Summary of the Invention by methods well known in the art. Some such procedures are described below.

25

- (i)       A compound of Formula (I) where R<sup>3</sup> is heterocyclalkyloxy can be prepared from a compound of formula 4 where Z is alkoxy as shown below:

- 45 -




A compound of Formula (I) where  $R^3$  is heterocyclalkyloxy can be prepared from a compound of formula 4 where Z is alkoxy by first de-alkylating the alkoxy group to give the corresponding compound of formula 5 where Z is hydroxy followed by reaction with a heterocyclalkyl halide [e.g., 4-(2-chloroethyl)morpholine, 1-(2-chloroethyl)pyrrolidine, and the like]. The de-alkylation reaction is carried out either with boron tribromide in a halogenated organic solvent such as dichloromethane or by heating 4 in neat pyridinium hydrochloride. The alkylation is carried out in the presence of a base (such as potassium carbonate, cesium carbonate, and the like) in a polar organic solvent such as acetonitrile, dimethylformamide, acetone, and the like.

Alternatively, a heterocyclalkyl group can be attached by reacting 5 with an alkyl dihalide followed by the reaction of the resulting haloalkyloxy intermediate with a heterocyclalkyl group (e.g., piperazine, morpholine, pyrrolidine, and the like) under the reaction conditions described above. Alkyl dihalides such as 1-bromo-2-chloroethane, 1-chloro-3-iodopropane, and the like, are commercially available.

(ii) A compound of Formula (I) where  $R^3$  is -O-(alkylene)- $R^9$  (where  $R^9$  is -COOH, -COR<sup>10</sup>, -COOR<sup>11</sup> or -CONR<sup>12</sup>R<sup>13</sup>) can be prepared from a compound of formula 5 as shown below:

- 46 -



- 5 A compound of Formula (I) where  $R^3$  is  $-O-(alkylene)-COOR^{11}$  is prepared by reacting a compound of formula **5** with an alkylating agent of formula  $X-(alkylene)-CO_2R^{11}$  where X is a halo group. Hydrolysis of the ester group provides the free acid ( $R^9$  is  $-COOH$ ) which can be converted to a compound of Formula (I) where  $R^9 = -CONR^{12}R^{13}$ , if desired, by treating the
- 10 acid with an amine of formula  $NR^{12}R^{13}$  (where  $R^{12}$  and  $R^{13}$  are as defined in the Summary of the Invention) in the presence of a suitable coupling agent (e.g., carbonyl diimidazole, N,N-dicyclohexylcarbodiimide and the like).

- 15 A compound of Formula (I) where  $R^9$  is  $-COR^{10}$  can be prepared from a compound of Formula (I) where  $R^9$  is  $-COOH$  by first converting the acid to a Weinreb amide followed by treatment with either a Grignard reagent or organolithium reagent of formula  $R^{10}MgBr$  or  $R^{10}Li$ , respectively.


- (iii) A compound of Formula (I) where  $R^3$  is  $-NH-(alkylene)-R^9$  where  $R^9$  is  $-COOH$ ,  $-COR^{10}$ ,  $-COOR^{11}$ ,  $-CONR^{12}R^{13}$  or heterocyclalkylamino can be
- 20 prepared from a compound of formula **4** where Z is a nitro group by reducing



- 47 -

the nitro group to the amino group and then following the procedures described above.

- (iv) A compound of Formula (I) where  $R^3$  is heteroalkenyl, heteroalkynyl, heterocyclyl-alkenyl, heterocyclylalkynyl, heteroalkyl or heterocyclylalkyl can be prepared as shown below.



- A compound of Formula (I) where  $R^3$  is heteroalkenyl, heteroalkynyl, heterocyclylalkenyl or heterocyclylalkynyl can be prepared by reacting a compound of formula 4 where Z is halo with a heteroalkene, heteroalkyne, heterocyclylalkene or heterocyclylalkyne respectively in the presence of a palladium (II) catalyst such as dichlorobis(triphenylphosphine)-palladium (II) in an organic base such as diisopropylamine, and the like.
- Heteroalkenes, heteroalkynes such as allyl alcohol, propargyl alcohol, 3-butyn-1-ol, propargylamine are commercially available. Heterocyclylalkyne can be prepared by reacting an alkynyl halide with a heterocycle. For example, 2-morpholin-1-ylprop-1-yne can be prepared by reacting propargyl bromide with morpholine. Reduction of the double or triple bond under catalytic hydrogenation reaction conditions provides the corresponding compound of Formula (I) where  $R^3$  is a heterocyclylalkyl or heteroalkyl group.

- (v) A compound of Formula (I) where  $R^3$  is  $-\text{NHSO}_2\text{R}^6$ ,  $-\text{NHSO}_2\text{NR}^7\text{R}^8$  or  $\text{NHC(X)R}^{23}\text{R}^{24}$  (where X is -O- or -S-) can be prepared from a compound of Formula (I) where  $R^3$  is amino by following the synthetic procedures described in PCT Application No. WO 97/46524.

- 48 -

A compound of Formula (I) where  $R^1$  is an acyl group can be prepared by reacting the the corresponding compound of Formula (I) where  $R^1$  is hydrogen with an acylating reagent of formula  $R^1COL$  where L is a leaving group under acylating reaction conditions such as halo. The reaction is  
5 carried out in the presence of a base such as sodium hydroxide, cesium carbonate, and the like.

#### Method (b)


10 Alternatively, a compound of Formula (I) can be prepared from an ester of formula 6 where Z is as defined above, by first converting the Z group in compound 6 to the desired  $R^3$  group utilizing the reaction conditions described in method (a)(i-v) above. Condensation of 7 with acetonitrile anion  
15 gives a 2-ketoacetonitrile of formula 8 which is then converted to a compound of Formula (I) utilizing the reaction conditions described in method (a) above.

Compounds of Formula (I) where  $R^2$  is thioalkyl or alkyl can be prepared by following the procedures described in U. S. Pat. No. 5,712,303.  
20

#### Scheme B

An alternate synthesis of compounds of Formula (I) where  $R^2$  is hydrogen and other groups are as defined in the Summary of the Invention  
25 is described below.

- 49 -




Condensation of 2-cyano-3-ethoxyacrylate of formula **9** with a hydrazine of formula **3** provides a 5-amino-4-ethoxycarbonyl pyrazole of formula **10**. The condensation reaction is carried out in a suitable polar organic solvent such as ethanol, isopropanol, and the like. Hydrolysis of **10** with an aqueous base (e.g., sodium hydroxide, lithium hydroxide, and the like) in an alcoholic organic solvent (e.g., methanol, ethanol, and the like) provides the corresponding 5-amino-4-carboxypyrazole of formula **11**.

Treatment of **11** with dipyridyldisulfide followed by reaction of the resulting thiopyridyl ester derivative **12** with an organometallic reagent such as a Grignard reagent or an organolithium reagent shown above provides a compound of Formula (I).

- 50 -

Scheme C

Another alternate synthesis of compounds of Formula (I) where  $R^2$  is hydrogen and other groups are as defined above is described below.



5

Thermal decarboxylation of a 5-amino-4-carboxypyrazole of formula **11** gives the corresponding 5-aminopyrazole of formula **13**. Compound **13**

- 51 -

is then converted to a compound of Formula (I) as shown in method (a) or (b) above.


In method (a), a compound of formula 13 is converted to a compound of Formula (I) by first protecting the amino group in compound 13 with a suitable amino protecting group (e.g., *tert*-butoxycarbonyl, and the like) to give the corresponding amino-protected compound of formula 14. Treatment of 14 with an acid derivative of formula  $R^3COL$  where L is a leaving group under organometallic displacement reaction conditions [e.g., alkoxy (preferably methoxy or ethoxy), dialkylamino, or preferably N,O-dimethylhydroxylamino] followed by the removal of the amino protecting group then provides a compound of Formula (I). The nucleophilic substitution is carried out in the presence of 2 equivalents of an alkylolithium (e.g., *tert*-butyllithium, and the like) and in an aprotic organic solvent such as tetrahydrofuran. The reaction conditions employed for the removal of the amino protecting group depends on the nature of the protecting group. For example, if *tert*-butoxycarbonyl is the protecting group, it is removed by treatment with an acid such as trifluoroacetic acid, hydrochloric acid, and the like.

Acid derivatives of formula  $R^3COL$  can be prepared by methods well known in the field of organic chemistry. For example, an acid derivative where L is a N,O-dimethylhydroxylamino group can be prepared from its corresponding acid by first converting the acid to the acid chloride with a suitable chlorinating agent such as oxalyl chloride, followed by treatment with N,O-dimethylhydroxylamine hydrochloride in the presence of an organic base such as triethylamine.

In method (b), a compound of formula 10 is brominated to give the 5-amino-4-bromo-pyrazole of formula 15. The bromination reaction is carried out with a suitable brominating agent such as N-bromosuccinimide in a suitable polar organic solvent such as dimethylformamide. Compound 15 is then converted to a compound of Formula (I) utilizing the reaction conditions described in Scheme C, method (a) above.

- 52 -

In a third aspect the present invention provides intermediates of formula



wherein Z is either hydroxy, nitro or halo group and A, B, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are as defined above.

In a fourth embodiment, the present invention provides medicaments or pharmaceutical compositions containing a compound of formula (I) and a pharmaceutically acceptable excipient.

### Utility

The compounds of Formula (I) are p38 MAP kinase inhibitors and therefore compounds of Formula (I) and compositions containing them are useful in the treatment of diseases such as rheumatoid arthritis, osteoarthritis, spondylitis, bone resorption diseases, sepsis, septic shock, toxic shock syndrome, endotoxic shock, tuberculosis, atherosclerosis, diabetes, adult respiratory distress syndrome, chronic pulmonary inflammatory disease, fever, periodontal diseases, ulcerative colitis, pyresis, Alzheimer's and Parkinson's diseases. Preferably, the compounds and medicaments may be used in the treatment of inflammatory diseases, preferably arthritis.

### Testing

The ability of the compounds of Formula (I) to inhibit p38 MAP kinase was demonstrated by the *in vitro* assay described in Example 15. The ability of the compounds of Formula (I) to inhibit the release of TNF- $\alpha$

- 53 -

was demonstrated by the *in vitro* and the *in vivo* assays described in detail in Examples 16 and 17, respectively. The anti-inflammatory activity of the compounds of this invention was determined utilizing adjuvant induced arthritis in rats assay described in Example 18.

5

### Administration and Pharmaceutical Compositions

In general, the compounds of this invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. The actual amount of the compound of this invention, i.e., the active ingredient, will depend upon numerous factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the route and form of administration, and other factors.

15

Therapeutically effective amounts of compounds of Formula (I) may range from approximately 0.1-50 mg per kilogram body weight of the recipient per day; preferably about 1-30 mg/kg/day. Thus, for administration to a 70 kg person, the dosage range would most preferably be about 70 mg to 2.1g per day.

20

In general, compounds of this invention will be administered as pharmaceutical compositions by any one of the following routes: oral, systemic (e.g., transdermal, intranasal or by suppository), or parenteral (e.g., intramuscular, intravenous or subcutaneous) administration. The preferred manner of administration is oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction. Compositions can take the form of tablets, pills, capsules, semisolids, powders, sustained release formulations, solutions, suspensions, elixirs, aerosols, or any other appropriate compositions.

30

The choice of formulation depends on various factors such as the mode of drug administration (e.g., for oral administration, formulations in the form of tablets, pills or capsules are preferred) and the bioavailability

- 54 -

of the drug substance. Recently, pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based upon the principle that bioavailability can be increased by increasing the surface area i.e., decreasing particle size. For example, U.S. Pat. No. 4,107,288  
5 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in which the active material is supported on a crosslinked matrix of macromolecules. U.S. Pat. No. 5,145,684 describes the production of a pharmaceutical formulation in which the drug  
10 substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to give a pharmaceutical formulation that exhibits remarkably high bioavailability.

The compositions are comprised of in general, a compound of  
15 Formula (I) in combination with at least one pharmaceutically acceptable excipient. Acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the compound of Formula (I). Such excipient may be any solid, liquid, semi-solid or, in the case of an aerosol composition, gaseous excipient that is generally available to one of  
20 skill in the art .

Solid pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium  
25 chloride, dried skim milk and the like. Liquid and semisolid excipients may be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, particularly for injectable solutions, include water, saline,  
30 aqueous dextrose, and glycols.

Compressed gases may be used to disperse a compound of this invention in aerosol form. Inert gases suitable for this purpose are nitrogen, carbon dioxide, etc.



- 55 -

Other suitable pharmaceutical excipients and their formulations are described in *Remington's Pharmaceutical Sciences*, edited by E. W. Martin (Mack Publishing Company, 18th ed., 1990).

5

The amount of the compound in a formulation can vary within the full range employed by those skilled in the art. Typically, the formulation will contain, on a weight percent (wt%) basis, from about 0.01-99.99 wt% of a compound of Formula (I) based on the total formulation, with the balance  
10 being one or more suitable pharmaceutical excipients. Preferably, the compound is present at a level of about 1-80 wt%. Representative pharmaceutical formulations containing a compound of Formula (I) are described in Example 14.


- 56 -

### EXAMPLES

The following preparations and examples are given to enable those skilled in the art to more clearly understand and to practice the present invention. They should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof. Numbers in brackets refer to the CPD # in Table I.

#### **Example 1**

5-Amino-1-(4-fluorophenyl)-4-[3-{3-(morpholin-4-yl)prop-1-ynyl}benzoyl]pyrazole (3)



#### Step 1

*n*-Butyllithium (214 ml, 340 mmol, 1.6 M solution in hexane) was added dropwise to a solution of acetonitrile (23.8 ml, 460 mmol) in dry tetrahydrofuran (1000 ml) at -78 °C. After stirring the reaction mixture for 20 min., a solution of 4-bromobenzoyl chloride in dry tetrahydrofuran (50 ml) was added dropwise over 20 min. After 1 h, saturated ammonium chloride was added (200 ml) and the reaction mixture was allowed to warm to room temperature. The product was extracted into ether and washed with 1N hydrochloric acid (400 ml). The organics were removed *in vacuo* and the residue was redissolved in ethyl acetate. Ammonium hydroxide was added to give a solid which was filtered, redissolved in ethyl acetate and washed with 2 N hydrochloric acid. The organic layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo* to give 2-(3-bromobenzoyl)-acetonitrile (16.6 g) as a solid.

#### Step 2

- 57 -

A mixture of 2-(3-bromobenzoyl)acetonitrile (16.5 g, 73.6 mmol) and N,N-diphenylformamidine (14.5 g, 73.6 mmol) in xylene (100 ml) was heated at reflux under a nitrogen atmosphere. After 3 h, the reaction mixture was cooled to room temperature and diluted with ether to give 2-(3-bromobenzoyl)-3-phenylaminoacrylonitrile (17.9 g) as a solid.

### Step 3

A mixture of 4-fluorophenylhydrazine (4.25 g, 33.7 mmol) and 2-(3-bromobenzoyl)-3-phenylaminoacrylonitrile (10.0 g, 30.7 mmol) in ethanol (100 ml) was heated at reflux under a nitrogen atmosphere. After 4 h, the reaction mixture was cooled to room temperature, diluted with hexane to give 5-amino-4-(3-bromobenzoyl)-1-(4-fluorophenyl)pyrazole (9.7 g) as a solid.

Replacing 4-fluorophenylhydrazine with 2,4-difluorophenylhydrazine in step 3 above gave 5-amino-4-(3-bromobenzoyl)-1-(2,4-difluorophenyl)pyrazole.

### Step 4

A mixture of 5-amino-4-(3-bromobenzoyl)-1-(4-fluorophenyl)pyrazole (1.3 g, 4.16 mmol), 4-(prop-2-ynyl)morpholine (2.1 g, 16.6 mmol) [prepared by adding to a solution of morpholine (14.7 ml, 168 mmol) in ether (50 ml), propargyl bromide (7.5 ml, 84 mmol) in ether (50 ml) dropwise over 30 min. and heating the reaction mixture to reflux. After 16 h, the reaction mixture was cooled to room temperature and filtered through a Buchner funnel. The filtrate was concentrated and purified by flash chromatography (gradient elution, 20-100% EtOAc/hexane) to give 4-(prop-2-ynyl)morpholine (5.0 g)], bis(triphenylphosphine)-palladium chloride (0.29 g, 0.42 mmol) and copper iodide (0.079 g, 0.42 mmol) in diisopropylamine (60 ml) was heated at 70 °C under argon. After 10 h, the reaction mixture was cooled to room temperature, diluted with ethyl acetate, washed with brine and dried over sodium sulfate. The organics were removed *in vacuo*. The crude product was purified by flash chromatography (elution gradient, EtOAc-5% MeOH/EtOAc with 0.2 % NH<sub>4</sub>OH) to give 5-amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylprop-1-ynyl)benzoyl]-pyrazole which was converted to the hydrochloride salt and

- 58 -

recrystallized from a mixture of methanol/ethyl acetate/hexane to give (1.4 g) of the pure product.


Proceeding as described in Example 1 above but substituting 4-(prop-2-ynyl)-morpholine in Step 4 with:

- 5 1-(prop-2-ynyl)-4-methylpiperazine,  
1-(prop-2-ynyl)piperidine,  
2-propyn-1-ol,  
1-dimethylamino-2-propyne, and  
2-methyl-3-butyn-2-ol; gave  
10 5-amino-1-(4-fluorophenyl)-4-[3-[3-(4-methylpiperazin-1-yl)prop-1-ynyl]benzoyl]pyrazole.2HCl (8),  
5-amino-1-(4-fluorophenyl)-4-[3-[3-(piperidin-1-yl)prop-1-ynyl]benzoyl]pyrazole.HCl (9),  
5-amino-1-(4-fluorophenyl)-4-[3-(3-hydroxyprop-1-ynyl)benzoyl]pyrazole  
15 (7),  
5-amino-4-[3-(3-dimethylaminoprop-1-ynyl)benzoyl]-1-(4-fluorophenyl)pyrazole.HCl (12), and  
5-amino-1-(4-fluorophenyl)-4-[3-(3-hydroxy-3-methyl-but-1-ynyl)benzoyl]pyrazole (87), respectively.

- 20 Proceeding as described in Example 1 above but substituting 4-fluorophenylhydrazine in Step 3 with 2,4-difluorophenylhydrazine, and 4-(prop-2-ynyl)morpholine in Step 4:  
with 3-ethynylpyridine gave  
5-amino-1-(2,4-difluorophenyl)-4-[3-(3-pyridylethynyl)benzoyl]pyrazole  
25 (88),  
with 3-(S,S-dioxo-thiomorpholin-4-yl)-1-propyne gave  
5-amino-1-(2,4-difluorophenyl)-4-[3-[3-(S,S-dioxo-thiomorpholin-4-yl)-1-propynyl]benzoyl]pyrazole (89), and  
with 1-ethynylcyclopentanol gave  
30 5-amino-1-(2,4-difluorophenyl)-4-[3-[2-(1-hydroxycyclopentyl)ethynyl]benzoyl]pyrazole (94).

**Example 2**

5-Amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylpropyl)benzoyl]-  
pyrazole hydrochloride (6)



5

A mixture of 5-amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylprop-  
1-ynyl)-benzoyl]pyrazole (0.45 g, 1.0 mmol) [prepared as described in  
Example 1] and 5 % Pd/C (0.07 g) in ethanol (20 ml) was stirred under  
10 hydrogen atmosphere. After 16 h, the reaction mixture was filtered  
through Celite® and the filtrate was concentrated *in vacuo*. The crude  
product was purified by flash chromatography (elution gradient, EtOAc-15  
% MeOH/EtOAc with 0.2 % NH<sub>4</sub>OH). The product was converted to the  
hydrochloride salt and recrystallized from a mixture of methanol/ethyl  
15 acetate to give 5-amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-  
ylpropyl)benzoyl]pyrazole.HCl (0.3 g, mpt. 211.9-212.6 °C) as a solid.

Proceeding as described in Example 2 above, but substituting of 5-  
amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylprop-1-  
ynyl)benzoyl]pyrazole with:  
20 5-amino-1-(4-fluorophenyl)-4-[3-[3-(4-methylpiperazin-1-yl)prop-1-  
ynyl]benzoyl]pyrazole,  
5-amino-1-(4-fluorophenyl)-4-[3-[3-(piperidin-1-yl)prop-1-  
ynyl]benzoyl]pyrazole,  
5-amino-1-(4-fluorophenyl)-4-[3-(3-hydroxyprop-1-ynyl)benzoyl]pyrazole,  
25 5-amino-4-[3-(3-dimethylaminoprop-1-ynyl)benzoyl]-1-(4-  
fluorophenyl)pyrazole,  
5-amino-1-(4-fluorophenyl)-4-[3-(3-hydroxy-3-methyl-1-  
butynyl)benzoyl]pyrazole,  
5-amino-1-(2,4-difluorophenyl)-4-[3-(3-pyridylethynyl)benzoyl]pyrazole,


- 60 -

5-amino-1-(2,4-difluorophenyl)-4-[3-{3-(S,S-dioxo-thiomorpholin-4-yl)-1-propynyl} benzoyl]pyrazole,  
5-amino-1-(4-fluorophenyl)-4-[3-{2-(1-hydroxycyclopentyl)ethynyl}benzoyl]pyrazole, and  
5 5-amino-1-(2,4-difluorophenyl)-4-[3-{2-(1-hydroxycyclopentyl)ethynyl}benzoyl]pyrazole gave  
5-amino-1-(4-fluorophenyl)-4-[3-[3-(4-methylpiperazin-1-yl)propyl]benzoyl]pyrazole (30);  
5-amino-1-(4-fluorophenyl)-4-[3-(3-piperidin-1-ylpropyl)benzoyl]pyrazole  
10 (32);  
5-amino-1-(4-fluorophenyl)-4-[3-(3-hydroxypropyl)benzoyl]pyrazole;  
5-amino-4-[3-(3-dimethylaminopropyl)benzoyl]-1-(4-fluorophenyl)pyrazole;  
5-amino-1-(4-fluorophenyl)-4-[3-(3-hydroxy-3-methylbutyl)benzoyl]pyrazole (90),  
15 5-amino-1-(2,4-difluorophenyl)-4-[3-(3-pyridylethyl)benzoyl]pyrazole (91),  
5-amino-1-(2,4-difluorophenyl)-4-[3-{3-(S,S-dioxo-thiomorpholin-4-yl)propyl} benzoyl]pyrazole (92),  
5-amino-1-(4-fluorophenyl)-4-[3-{2-(1-hydroxycyclopentyl)ethyl}benzoyl]pyrazole (93), and  
20 5-amino-1-(2,4-difluorophenyl)-4-[3-{2-(1-hydroxycyclopentyl)ethyl}benzoyl]pyrazole (94) respectively.

- 61 -

**Example 3**

5-Amino-1-(4-fluorophenyl)-4-[3-{2-(morpholin-4-yl)ethoxy}benzoyl]-pyrazole hydrochloride (14)

**Step 1**

A mixture of methyl 3-hydroxybenzoate (8.0 g, 56 mmol) and 4-(2-chloroethyl)-morpholine hydrochloride (15.7 g, 84 mmol) and potassium carbonate (11.5 g, 83 mmol) in toluene (50 ml) was heated at reflux. After 4 days, the reaction mixture was cooled to room temperature and diluted with ethyl acetate. The organic layer was washed with water and then extracted with dilute hydrochloric acid. The acidic layer was separated, basified with 5 N sodium hydroxide and the product was extracted into ethyl acetate. The organics were removed *in vacuo* and the residue was purified by flash chromatography (elution gradient 3 % acetone/methylene chloride) to give methyl 3-(2-morpholin-4-ylethoxy)benzoate (9.0 g) as an oil.

**Step 2**

Lithium diisopropylamide (18.8 ml, 37 mmol, 2.0 M solution in heptane/ tetrahydrofuran/ethylbenzene) was added dropwise to a solution of acetonitrile (1.58 g, 37 mmol) in dry tetrahydrofuran (50 ml) at -78 °C. After stirring the reaction mixture for 30 min., a solution of methyl 3-(2-morpholin-4-ylethoxy)benzoate in dry tetrahydrofuran (50 ml) was added dropwise over 10 min. After 15 min., water was added and the reaction mixture was allowed to warm to room temperature. The aqueous layer was separated and acidified with dilute hydrochloric acid to pH 7. The product was extracted into ethyl acetate and washed with water and brine and dried over magnesium sulfate. The organics were removed *in vacuo* to

- 62 -

give 2-[3-(2-morpholin-4-ylethoxy)phenyl]acetonitrile (5.0 g) as an oil which was used in the next step without further purification.

### Step 3

A mixture of 2-[3-(2-morpholin-4-ylethoxy)phenyl]acetonitrile (5.0 g) and N,N-diphenylformamidine (5.0 g, 25.5 mmol) in xylene (150 ml) was heated at 100 °C under a nitrogen atmosphere. After 3 h, the reaction mixture was cooled to room temperature and diluted with hexane to give 2-[3-(2-morpholin-4-ylethoxy)benzoyl]-3-phenylamino-acrylonitrile (5.0 g) as a solid.

### Step 4


A mixture of 4-fluorophenylhydrazine (1.0 g, 6.8 mmol) and 2-[3-(2-morpholin-4-ylethoxy)-benzoyl]-3-phenylaminoacrylonitrile (2.0 g, 5.3 mmol) in ethanol (30 ml) was heated at reflux under a nitrogen atmosphere. After 6 h, the reaction mixture was cooled to room temperature and diluted with water. The product was extracted into ethyl acetate and the organic layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. Purification by flash chromatography (elution gradient: CH<sub>2</sub>Cl<sub>2</sub>-3%MeOH/CH<sub>2</sub>Cl<sub>2</sub>) gave 5-amino-1-(4-fluorophenyl)-4-[3-(2-morpholin-4-ylethoxy)benzoyl]pyrazole which was converted to the hydrochloride salt (0.7 g, mpt. 191.6-192.5 °C).

Replacing 4-fluorophenylhydrazine in Step 4 above with:  
2-fluorophenylhydrazine, and  
2,6-dichlorophenylhydrazine, respectively were obtained:  
5-amino-1-(2-fluorophenyl)-4-[3-(2-morpholin-4-ylethoxy)benzoyl]pyrazole (97), and  
5-amino-1-(2,6-dichlorophenyl)-4-[3-(2-morpholin-4-ylethoxy)benzoyl]pyrazole (98).



### Example 4

#### 5-Amino-1-(4-fluorophenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (20)



##### Step 1

A mixture of 5-amino-4-(3-bromobenzoyl)-1-(4-fluorophenyl)pyrazole (0.9 g, 2.5 mmol) [prepared as described in Example 1 above], pyridine-3-boronic acid, 1,3-propanediol cyclic ester (0.5 g, 3 mmol), potassium phosphate (0.8 g, 3.73 mmol) and tetrakis(triphenylphosphine) palladium (0.3 g, 0.25 mmol) in dioxane (20 ml) was heated at 85 °C under argon. After 16 h, the reaction mixture was cooled to room temperature and poured into brine. The product was extracted into ethyl acetate, dried over sodium sulfate and filtered. The organic layer was removed *in vacuo* and the residue was purified by flash chromatography (elution gradient: 40-80% ethyl acetate/hexane) to give 5-amino-1-(4-fluorophenyl)-4-[3-(pyridin-3-yl)benzoyl]-pyrazole (0.50 g) which was recrystallized from ethyl acetate (mpt. 222.2-223.0).


Treatment of 5-amino-1-(4-fluorophenyl)-4-[3-(pyridin-3-yl)benzoyl]-pyrazole with methyl iodide in ethyl acetate gave 5-amino-1-(4-fluorophenyl)-4-[3-(N-methylpyridinium-3-yl)benzoyl]pyrazole iodide (59).

Substitution of 5-amino-4-(3-bromobenzoyl)-1-(4-fluorophenyl)pyrazole with 5-amino-4-(3-bromobenzoyl)-1-(2,4-difluorophenyl)pyrazole in Step 1 above followed by conversion to the hydrochloride salt gave 5-amino-1-(2,4-difluorophenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole.HCl (99).

- 64 -

**Example 5**

5-Amino-4-[3-(2-aminosulfonylethyl)benzoyl]-1-(4-fluorophenyl)pyrazole(50)

**Step 1**

A mixture of 5-amino-4-(3-bromobenzoyl)-1-(4-fluorophenyl)pyrazole (1.5 g, 4.14 mmol) [prepared as described in Example 1 above], vinylsulfonamide (1.33 g, 12.4 mmol), bis(triphenylphosphine)palladium chloride (0.3 g, 0.42 mmol) and triethylamine (6 ml, 43 mmol) in dimethylformamide (18 ml) was heated at 100 °C under argon. After 16 h, the reaction mixture was cooled to room temperature and poured into 1 N hydrochloric acid. The product was extracted into ethyl acetate, washed with brine, dried over sodium sulfate and filtered. The organic layer was removed *in vacuo* and the residue was purified by flash chromatography (elution gradient: 40-80% ethyl acetate/hexane) to give 5-amino-4-[3-(2-aminosulfonylethenyl)benzoyl]-1-(4-fluorophenyl)pyrazole which was recrystallized from a mixture of methanol/ethyl acetate/hexane to give 0.78 g of the desired product.

**Step 2**

A mixture of 5-amino-4-[3-(2-aminosulfonylethenyl)benzoyl]-1-(4-fluorophenyl)-pyrazole (2.1 g, 5.43 mmol) and palladium hydroxide (0.6 g) in methanol (150 ml) was shaken in a Parr apparatus under hydrogen atmosphere at 50 psi. After 4 days, the reaction mixture was filtered through Celite® and the filtrate was concentrated. The residue was purified by flash chromatography (elution gradient: 40-100% ethyl acetate/hexane) to give a crude product which was recrystallized from


- 65 -

methanol/ethyl acetate/hexane to give 5-amino-4-[3-(2-aminosulfonylethyl)benzoyl]-1-(4-fluorophenyl)pyrazole (0.95 g, mpt. 170-170.4 °C) as a solid.

Replacement of vinylsulfonamide in Step 1 above with vinylmethylsulfone gave:  
5-amino-4-[3-(2-methylsulfonylethyl)benzoyl]-1-(4-fluorophenyl)pyrazole (100).

### Example 6

5-Amino-1-(4-fluorophenyl)-4-[3-(morpholin-4-ylmethylcarbonyl)benzoyl]pyrazole(18)



#### Step 1

A mixture of 5-amino-4-(3-bromobenzoyl)-1-(4-fluorophenyl)pyrazole (3.5 g, 9.7 mmol) [prepared as described in Example 1 above], tributyl-(1-ethoxyvinyl)tin (4.3 ml, 12.36 mmol) and tetrakis(triphenylphosphine)palladium (1.0 g, 0.87 mmol) in dimethylformamide (25 ml) was heated at 95 °C under argon. After 16 h, the reaction mixture was cooled to room temperature and 10% aqueous hydrochloric acid (25 ml) was slowly added. After 30 min., the reaction mixture was diluted with ethyl acetate and filtered through Celite®. The organic layer was separated and washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: 10-60% ethyl acetate/hexane) to give 5-amino-4-[3-(1-ethoxyvinyl)benzoyl]-1-(4-fluorophenyl)pyrazole which was dissolved in tetrahydrofuran (50 ml). 1 N hydrochloric acid (20 ml) was added and the reaction mixture was stirred at room temperature for 16 h.

- 66 -

The organic layer was separated, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The crude product was purified by flash chromatography (elution gradient: 20-50% ethyl acetate/hexane) and then was recrystallized from a mixture of ethyl acetate/hexane to give 5-amino-4-[3-acetylbenzoyl]-1-(4-fluorophenyl)pyrazole (2.0 g).

### Step 2

To a suspension of copper bromide (2.2 g, 9.85 mmol) in a (1:1) mixture of ethyl acetate/methylene chloride (100 ml) at reflux was added a solution of 5-amino-4-[3-acetylbenzoyl]-1-(4-fluorophenyl)pyrazole (1.6 g, 4.95 mmol) in methylene chloride (25 ml) under nitrogen. After 16 h, the reaction mixture was concentrated and the residue was partitioned between aqueous sodium bisulfite and ethyl acetate. The organic layer was separated, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: 10-40% ethyl acetate/hexane) to give 5-amino-4-[3-(2-bromoacetyl)benzoyl]-1-(4-fluorophenyl)pyrazole (0.47 g) as a solid.


### Step 3

To a solution of morpholine (0.25 ml, 2.79 mmol) in dimethylformamide (5 ml) was added a solution of 5-amino-4-[3-(2-bromoacetyl)benzoyl]-1-(4-fluorophenyl)pyrazole (0.22 g, 0.56 mmol) in dimethylformamide (5 ml). After 16 h, the reaction mixture was poured into brine and the product was extracted into ethyl acetate. The organic layer was separated, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: ethyl acetate-10% methanol/ethyl acetate) to give 5-amino-1-(4-fluorophenyl)-4-[3-(morpholin-4-ylmethylcarbonyl)benzoyl]pyrazole (0.05 g) as a solid.

- 67 -

**Example 7**

5-amino-1-(4-fluorophenyl)- 4-[3-(2-hydroxyethyl)benzoyl]pyrazole (118)

**Step 1**

To a solution of 3-bromophenylacetic acid (10 g, 46.5 mmol) in tetrahydrofuran (100 ml) at 0 °C was added diborane (70 ml, 1.0 M solution in tetrahydrofuran). The reaction mixture was allowed to warm to room temperature. After 16 h, the reaction mixture was cooled to 0 °C and water was added dropwise (50 ml). The organic layer was separated and washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: 40-60% ethyl acetate/hexane) to give 3-(2-hydroxyethyl)bromobenzene (9.0 g).

**Step 2**

To a solution of 3-(2-hydroxyethyl)bromobenzene (4.0 g, 20 mmol) in methylene chloride (100 ml) at 0 °C was added a solution of *tert*-butyldimethylsilyl chloride (3.6 g, 24 mmol), dimethylaminopyridine (0.61 g, 5 mmol) and triethylamine (3.6 ml, 25.9 mmol). After 1 h, the reaction mixture was washed with brine, saturated ammonium chloride, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: 0-10% hexane/ethyl acetate) to give 3-(2-*tert*-butyl-dimethylsiloxyethyl)bromobenzene (6.0 g).

**Step 3**

A mixture of ethyl (ethoxymethylene)cynoacetate (26 ml, 154 mmol) and 4-fluorophenyl hydrazine (19.4 g, 154 mmol) in ethanol (125 ml) was heated at reflux. After 16 h, the reaction mixture was cooled to room temperature. The solid was filtered off and dried to give 5-amino-4-ethylcarboxy-1-(4-fluorophenyl)pyrazole (28 g) which was suspended in a

- 68 -

mixture of 1 N lithium hydroxide (100 ml) and methanol (250 ml). The reaction mixture was heated at reflux. After 16 h, the reaction mixture was filtered through a sinter funnel and the filtrate was acidified with 2 N hydrochloric acid (65 ml). The solid was filtered off and dried to give 5-amino-4-carboxy-1-(4-fluorophenyl)pyrazole (21 g).

#### Step 4

A mixture of 5-amino-4-carboxy-1-(4-fluorophenyl)pyrazole (15 g, 68 mmol), aldrathiol-2 (14.9 g, 68 mmol) and triphenylphosphine (17.8 g, 68 mmol) in acetonitrile (2 l) was stirred at room temperature. After 16 h, the product was filtered off and dried to give 5-amino-1-(4-fluorophenyl)-4-(2-pyridylthiocarboxy)pyrazole (14 g).

#### Step 5

Into an oven dried flask containing magnesium turnings (0.386 g, 15.9 mmol) and tetrahydrofuran (10 ml) was added 3-(2-*tert*-butyldimethylsiloxyethyl)bromobenzene (5.0 g, 15.9 mmol) and the reaction mixture was heated at reflux. After 3 h, the reaction mixture was cooled to room temperature and 5-amino-1-(4-fluorophenyl)-4-(2-pyridylthiocarboxy)pyrazole (2.37 g, 7.6 mmol) was added and the stirring was continued for 16 h. The reaction mixture was concentrated *in vacuo*. The residue was dissolved in ethyl acetate and washed with aqueous ammonium chloride and brine and dried over sodium sulfate. The organics were removed *in vacuo* and the residue was purified by flash chromatography (elution gradient: 10-30% ethyl acetate/hexane) to give 5-amino-1-(4-fluorophenyl)-4-[3-(2-*tert*-butyldimethylsiloxyethyl)benzoyl]pyrazole (1.20 g).

#### Step 6

To a solution of 5-amino-1-(4-fluorophenyl)-4-[3-(2-*tert*-butyldimethylsiloxyethyl)benzoyl]pyrazole (1.2 g, 3.0 mmol) in tetrahydrofuran (25 ml) was added tetrabutyl-ammonium fluoride (3.6 ml, 3.6 mmol, 1 M solution in tetrahydrofuran). After 1 h, the reaction mixture was poured into brine and the product was extracted into ethyl acetate. The organic layer was dried over sodium sulfate, filtered and concentrated *in vacuo*. The residue was purified by flash chromatography


- 69 -

(elution gradient: 40-100 % ethyl acetate/hexane) to give 5-amino-1-(4-fluorophenyl)-4-[3-(2-hydroxyethyl)benzoyl]pyrazole (0.8 g).

5

### Example 8

Synthesis of 5-amino-1-(4-fluorophenyl)-4-[3-[4-methylpiperazin-1-yl)ethyl)-benzoyl]pyrazole dihydrochloride (31)



10

#### Step 1

To a solution of 5-amino-1-(4-fluorophenyl)-4-[3-(2-hydroxyethyl)benzoyl]pyrazole (0.8 g, 2.5 mmol) in pyridine (10 ml) was added methanesulfonyl chloride (0.29 ml, 3.7 mmol). After 2 h, the reaction mixture was poured into 2 N hydrochloric acid (40 ml) and the product was extracted into ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: 40-100 % ethyl acetate/hexane) to give 5-amino-1-(4-fluorophenyl)-4-[3-(2-methanesulfonyloxyethyl)benzoyl]pyrazole (0.87 g).

20

#### Step 2

A mixture of 5-amino-1-(4-fluorophenyl)-4-[3-(2-methanesulfonyloxyethyl)-benzoyl]pyrazole (0.22 g, 0.55 mmol), N-methylpiperazine (0.18 ml, 1.64 mmol) and potassium carbonate (0.22 g, 1.64 mmol) in dimethylformamide (10 ml) was heated at 70 °C. After 4 h, the reaction mixture was cooled to room temperature, poured into water and the product was extracted into ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo*. The residue was purified by flash chromatography (elution

25


- 70 -

gradient: ethyl acetate- 20% methanol/ethyl acetate) to 5-amino-1-(4-fluorophenyl)-4-[3-[4-methylpiperazin-1-yl)ethyl)benzoyl]pyrazole which was converted to the hydrochloride salt (mpt. 272.9-273.9).

5

**Example 9**

5-Amino-4-[3-(2-aminoethyl)benzoyl]-1-(4-fluorophenyl)-pyrazole hydrochloride (47)



10

**Step 1**

A mixture of 5-amino-1-(4-fluorophenyl)-4-[3-(2-methanesulfonyloxyethyl)-benzoyl]pyrazole (0.40 g, 0.99 mmol), sodium azide (0.19 ml, 2.97 mmol) and potassium carbonate (0.41 g, 2.97 mmol) in dimethylformamide (15 ml) was stirred at room temperature. After 16 h, the reaction mixture was poured into brine and the product was extracted into ethyl acetate. The organic layer was dried over sodium sulfate, filtered and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: 20-50 % ethyl acetate/hexane) to 5-amino-1-(4-fluorophenyl)-4-[3-(2-azidoethyl)benzoyl]pyrazole (0.32 g).

20

**Step 2**

To a solution of 5-amino-1-(4-fluorophenyl)-4-[3-(2-azidoethyl)benzoyl]pyrazole (0.31 g, 0.9 mmol) in tetrahydrofuran (15 ml) was added triphenylphosphine (3.55 g, 1.36 mmol). After 48 h, the reaction mixture concentrated *in vacuo*. The residue was dissolved in 2 N sodium hydroxide and the product was extracted into ethyl acetate. The organic layer was dried over sodium sulfate, filtered and concentrated *in vacuo*. The product was converted to its hydrochloride salt and recrystallized from

25




- 71 -

a mixture of methanol-ethyl acetate to give 5-amino-4-[3-(2-aminoethyl)benzoyl]-1-(4-fluorophenyl)pyrazole hydrochloride salt (0.22 g).

### Example 10

#### 5-Amino-4-[3-(*tert*-butoxycarbonylmethoxy)benzoyl]-1-(4-fluorophenyl)pyrazole



#### Step 1

Into an oven dried flask containing magnesium turnings (0.408 g, 17 mmol) and tetrahydrofuran (10 ml) was added 3-bromoanisole (3.1 g, 17 mmol) and the reaction mixture was heated at reflux. After 2 h, the reaction mixture was cooled to room temperature and 5-amino-1-(4-fluorophenyl)-4-(2-pyridylthiocarboxy)pyrazole (1.5g, 4.8 mmol) was added and the stirring was continued for 1 h. The reaction mixture was quenched with water and the product was extracted into ethyl acetate. The organic layer was washed with aqueous ammonium chloride and brine and dried over sodium sulfate. The organics were removed *in vacuo* and the residue was filtered and washed with hexane to give 5-amino-1-(4-fluoro-phenyl)-4-(3-methoxybenzoyl)pyrazole (1.20 g).

#### Step 2

To an ice-cooled solution of 5-amino-1-(4-fluorophenyl)-4-(3-methoxybenzoyl)-pyrazole (3.0 g, 10.0 mmol) in methylene chloride (25 ml) was added boron tribromide (51 ml, 51 mmol, 1 M solution in methylene chloride). After 1 h, the reaction mixture was poured into brine and the product was extracted into ethyl acetate. The organic layer was dried over

- 72 -


sodium sulfate, filtered and concentrated *in vacuo* to give 5-amino-1-(4-fluorophenyl)-4-[3-hydroxybenzoyl]pyrazole (2.4 g).

### Step 3

A mixture of 5-amino-1-(4-fluorophenyl)-4-[3-hydroxybenzoyl]pyrazole (1.0 g, 3.3 mmol), *tert*-butyl bromoacetate (1.4 g, 7.2 mmol) and potassium carbonate (1 g, 7.2 mmol) in acetonitrile was heated at 70 °C overnight. The reaction mixture was cooled, diluted with ethyl acetate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash chromatography (elution gradient: 10% acetone/hexane) to give 5-amino-4-[3-(*tert*-butoxycarbonylmethoxy)benzoyl]-1-(4-fluorophenyl)pyrazole (1.2 g) as a solid.


### Example 11

5-Amino-4-[3-carboxymethoxybenzoyl]-1-(4-fluorophenyl)-pyrazole (119)



### Step 1

A mixture of 5-amino-4-[3-(*tert*-butoxycarbonylmethoxy)benzoyl]-1-(4-fluorophenyl)pyrazole (1.0 g, 3.3 mmol) and trifluoroacetic acid (15 ml, 194 mmol) in methylene chloride (15 ml) was stirred overnight at room temperature. The organics were removed *in vacuo* and the residue was dissolved in toluene. The solution was concentrated and the residue was triturated between ethyl acetate and hexane to give 5-amino-4-[3-(carboxymethoxy)benzoyl]-1-(4-fluorophenyl)pyrazole (0.8 g) as a solid.

**Example 12****5-Amino-1-(4-fluorophenyl)-4-[3-(methylaminocarbonylmethoxy)-benzoyl]pyrazole (34)****Step1**


To a solution of 5-amino-4-[3-(carboxymethoxy)benzoyl]-1-(4-fluorophenyl)-pyrazole (0.5 g, 1.43 mmol) in tetrahydrofuran (10 ml) was added carbonyl diimidazole (0.3 g, 1.85 mmol) and the reaction mixture was heated at 60 °C. After 1 h, methylamine (10 ml, 5 mmol, 0.5 M solution in tetrahydrofuran) was added and reaction was continued at 60 °C overnight. The reaction mixture was cooled and diluted with ethyl acetate. The organic layer was separated and washed with brine and dried over sodium sulfate. The organics were removed *in vacuo* and the residue was purified by flash chromatography (elution gradient: 20-30% acetone/hexane) to give 5-amino-1-(4-fluorophenyl)-4-[3-(methylaminocarbonyl-methoxy)benzoyl]pyrazole (0.25 g, mpt. 195.6-196.3 °C) as a solid.

Proceeding as described in Example 12 above, but substituting methylamine with: morpholine gave 5-amino-1-(4-fluorophenyl)-4-[3-(morpholin-4-ylcarbonylmethoxy)-benzoyl]pyrazole (35).

**Example 13**

5-amino-1-(4-fluorophenyl)-4-[3-{3-(morpholin-4-yl)propylamino}benzoyl]-pyrazole (48)

5

**Step 1**

Benzoylacetonitrile (14.5 g, 10 mmol) was added to cold fuming nitric acid (50 ml) portionwise over 10 min. The reaction mixture was stirred for 15 min., and then poured into ice. The solid was filtered off and recrystallised from ethanol to give 2-(3-nitrobenzoyl)-acetonitrile (5.4 g) as a brown solid.

**Step 2**

A mixture of 2-(3-nitrobenzoyl)acetonitrile (13.75 g, 72.3 mmol) and N,N-diphenylformamidine (14.2 g, 72.3 mmol) in xylene (200 ml) was heated at reflux under nitrogen atmosphere. After 3 h, the reaction mixture was cooled to room temperature and diluted with xylenes to give 2-(3-nitrobenzoyl)-3-phenylaminoacrylonitrile (15.7 g) as a yellow solid.

**Step 3**

A mixture of 4-fluorophenylhydrazine (2.24 g, 15.57 mmol) and 2-(3-nitrobenzoyl)-3-phenylaminoacrylonitrile (4.15 g, 14.16 mmol) in ethanol (50 ml) was heated at reflux under nitrogen atmosphere. After 1 h, the reaction mixture was cooled to room temperature and stirred for an additional 3 h. The solid was filtered and dried to give 5-amino-1-(4-fluorophenyl)-4-(3-nitrobenzoyl)pyrazole (4.5 g) as a solid.

**Step 4**

A mixture of 5-amino-1-(4-fluorophenyl)-4-(3-nitrobenzoyl)pyrazole (4.0 g, 24.52), Fe powder (3.84 g, 68 mmol) and ammonium chloride (3.84, 71.78 mmol) in ethanol (135 ml) and water (64 ml) was heated at reflux

- 75 -

under nitrogen atmosphere. After 1 h, the reaction mixture was cooled to room temperature and stirred overnight. The reaction mixture was filtered through Celite® and the filtrate was concentrated *in vacuo*. The residue was partitioned between water and ethyl acetate. The organic layer was separated and washed with brine, dried over sodium sulfate and concentrated *in vacuo* to give 5-amino-4-(3-amino-benzoyl)-1-(4-fluorophenyl)pyrazole (3.5 g) as a solid.

#### Step 5

5-amino-4-(3-aminobenzoyl)-1-(4-fluorophenyl)pyrazole (0.5 g, 1.6 mmol), 1-bromo-3-chloropropane (0.26 g, 1.6 mmol) and cesium carbonate (0.52 g, 1.6 mmol) in dimethylformamide (25 ml) was heated at 80 °C. After 2 days, the reaction mixture was cooled to room temperature and diluted with ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: 20 % acetone/hexanes) to give 5-amino-4-[3-(3-chloropropylamino)benzoyl]-1-(4-fluorophenyl)pyrazole (0.2 g) as a solid.

#### Step 6


A mixture of 5-amino-4-[3-(3-chloropropylamino)benzoyl]-1-(4-fluorophenyl)-pyrazole (0.05 g, 0.13 mmol), morpholine (0.1 ml, 1.1 mmol), potassium carbonate (0.1 g) and potassium iodide (0.1 g) in acetonitrile (3 ml) was heated at reflux. After 2 days, the reaction mixture was poured into brine and the product was extracted into ethyl acetate. The organic layer was separated, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash chromatography (elution gradient: 3 % MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give 5-amino-1-(4-fluorophenyl)-4-[3-{3-(morpholin-4-yl)propylamino}-benzoyl]pyrazole as a solid.

- 76 -

**Example 14**

5-amino-1-(4-fluorophenyl)-4-[3-{2-(piperidin-1-yl)ethoxy}benzoyl]pyrazole  
HCl salt (81)

5

**Step 1**

5-Amino-1-(4-fluorophenyl)-4-[3-hydroxybenzoyl]pyrazole, from  
10 Example 10, step 2, 1.5g, 5.05mmol) was combined with toluene (50mL).  
2-bromoethanol (1.79 mL, 25.23 mmol) was added and then the reaction  
mixture was cooled to 0°C. Triphenylphosphine (5.425g, 20.69mmol) and  
diethyl azodicarboxylate (3.26 mL, 20.69 mmol) were then added. The  
reaction was allowed to warm to room temperature. After stirring for 16  
15 hours, the reaction was quenched with a saturated aqueous solution of  
NH<sub>4</sub>Cl, extracted with ethyl acetate, dried (MgSO<sub>4</sub>), filtered, and  
concentrated under vacuum. The product (5-amino-1-(4-fluorophenyl)-4-[3-  
(2-bromoethoxy)benzoyl]pyrazole) was purified by column chromatography  
on silica gel using 40:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH then stirred with ether for 20  
20 minutes, filtered and dried to give 0.785g of product.

**Step 2**

5-amino-1-(4-fluorophenyl)-4-[3-(2-bromoethoxy)benzoyl]pyrazole  
(0.6g, 1.48 mmol) was combined with piperidine (1.47 mL, 14.8 mmol) and  
ethanol (10mL) and heated at reflux for 16 hrs. The reaction mixture was  
25 concentrated under vacuum. The resulting residue was partitioned  
between a saturated aqueous solution of NaHCO<sub>3</sub> and ethyl acetate and  
extracted three times with ethyl acetate. The organic extracts were dried  
(MgSO<sub>4</sub>), filtered, concentrated under vacuum and purified by column




- 77 -

chromatography on silica gel using 16:1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH. Dissolving the product in ethyl acetate then adding hydrochloric acid (1.0M, 1.0 equivalent) formed the hydrochloric salt which was filtered and dried to give 0.413g of 5-amino-1-(4-fluorophenyl)-4-[3-{2-(piperidin-1-yl)ethoxy}benzoyl]pyrazole.HCl (mpt 210.2-211.2°C).




Proceeding as in Step 2 but replacing piperidine with diethanolamine, dimethylamine, N-methylpiperazine, 2-aminoethanol, bis(2-methoxyethyl)amine, diethylamine, methylamine, ammonia, and 3-oxopyridazine the following compounds were obtained.

10




- 78 -

Mole structure	CPD#	HRMS MW	Melting Point
 ClH	121	368.41	184.5-190
 ClH	39	464.922	160.3-160.8
 ClH	122	494.95	238.0-258.0




 <p>Chemical structure of compound 123: 1-(4-fluorophenyl)-4-((4-(4-((2-((4-chlorophenyl)oxy)ethyl)amino)ethyl)phenyl)-2-oxo-1-phenylethyl)-1H-1,2,4-triazol-5-amine. The structure shows a triazole ring with an amino group at position 5, a 4-fluorophenyl group at position 1, and a 2-oxo-1-phenylethyl group at position 4. The ketone oxygen is double-bonded to the ethyl chain. The ethyl chain is connected via an ether linkage to a 4-(4-((2-((4-chlorophenyl)oxy)ethyl)amino)ethyl)phenyl group.</p> <p>ClH</p>	123	425.4	217.1-218.0
 <p>Chemical structure of compound 124: 1-(4-fluorophenyl)-4-((4-(4-((2-((4-(methoxymethoxy)ethyl)amino)ethyl)phenyl)-2-oxo-1-phenylethyl)-1H-1,2,4-triazol-5-amine. The structure is similar to compound 123, but the 4-chlorophenyl group is replaced by a 4-(methoxymethoxy)ethyl group, represented as H<sub>3</sub>C-O-CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>3</sub>.</p> <p>ClH</p>	124	506.49	86.5-93.5
 <p>Chemical structure of compound 125: 1-(4-fluorophenyl)-4-((4-(4-((2-diethylamino)ethyl)phenyl)-2-oxo-1-phenylethyl)-1H-1,2,4-triazol-5-amine. The structure is similar to compound 123, but the 4-chlorophenyl group is replaced by a 4-diethylaminoethyl group, represented as -CH<sub>2</sub>-CH<sub>2</sub>-N(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>.</p> <p>ClH</p>	125	440.13	137.8-139.8

- 80 -

 ClH	126	396.25	204.9-210.3
 ClH	127	376.817	231.5-232.5
	128	419.414	174.5-178.0

**Example 15**

5-amino-1-(4-fluorophenyl)-4-[3-(pyridin-2-ylmethoxy)benzoyl]pyrazole (82)



5

5-Amino-1-(4-fluorophenyl)-4-[3-hydroxybenzoyl]pyrazole, from Example 10, step 2, (0.5g, 1.68 mmol), 2-pyridylcarbinol (0.81 mL, 8.41 mmol), triphenylphosphine (1.81g, 6.9 mmol), and diethylazodicarboxylate (1.09 mL, 6.9 mmol) were combined in toluene (50mL). The reaction mixture was stirred for 16 hours then quenched with a saturated aqueous solution of  $\text{NH}_4\text{Cl}$  and extracted three times with ethyl acetate. The product was then extracted from the ethyl acetate into a 10% aqueous solution of HCl. The aqueous layer was then neutralized with NaOH and extracted with ethyl acetate. The organic extracts were dried ( $\text{MgSO}_4$ ), filtered, and concentrated under vacuum. The residue was purified by column chromatography on silica gel using 1:1 hexane/ethyl acetate to give 0.165g of 5-amino-1-(4-fluorophenyl)-4-[3-(pyridin-2-ylmethoxy)benzoyl]pyrazole (mpt. 176.1-177.3°C).

20


Replacing 2-pyridylcarbinol with glycolic acid, 1-(2-hydroxyethyl)-2-pyrrolidinone and 4-hydroxypiperidine gave the following compounds.

- 82 -

Mole structure	CPD#	HRMS MWt	M. Pt.
	119	355.324	215.9-216.2
	129	412.03	
	130	416.882	195.0-220.0


**Example 16**

5-amino-1-(4-fluorophenyl)-4-[3-isopropylaminocarbonyloxybenzoyl]pyrazole (83)



5-Amino-1-(4-fluorophenyl)-4-[3-hydroxybenzoyl]pyrazole, from Example 10, step 2, (0.30g, 1.01 mmol) was combined with  $K_2CO_3$  (0.418g, 3.03 mmol) and THF (6mL). The mixture was cooled in an ice bath and then isopropyl isocyanate (0.12 mL, 1.21 mmol) was added. The reaction was allowed to warm to room temperature and stirred for 16 hours. The reaction mixture was quenched with water, extracted into ethyl acetate, dried ( $MgSO_4$ ), filtered, and concentrated to dryness. The residue was stirred in methanol and dichloromethane for one hour then filtered to give 0.118g of 5-amino-1-(4-fluorophenyl)-4-[3-isopropylaminocarbonyloxybenzoyl]pyrazole (mpt. 225.2-230.1°C).

Replacing isopropyl isocyanate with ethyl isocyanate was made 5-amino-1-(4-fluorophenyl)-4-[3-ethylaminocarbonyloxybenzoyl]pyrazole (84). Mpt. 201.2-202.8°C.

**Example 17****5-Amino-1-(4-fluorophenyl)-4-[3-iodo  
benzoyl]pyrazole**5 **Step 1**

*n*-Butyllithium (30.5 ml, 76 mmol, 2.5 M solution in hexane) was added dropwise to a cooled (0 °C) solution of diisopropylamine (10.6 ml, 76 mmol) in 10 ml dry tetrahydrofuran. Once addition was complete, the solution was kept at 0 °C for 10 minutes and was then cooled to -50 °C.

- 10 This cold LDA solution was then added to a -50 °C solution of acetonitrile (2.37 ml, 45.3 mmol) and ethyl 4-iodobenzoate (10.0g, 36.2 mmol) in dry tetrahydrofuran (18 ml). Once addition was complete, the reaction was stirred at -50 °C for 3 hours and was subsequently warmed to 0 °C. Saturated ammonium chloride was added (20 ml) and the reaction mixture
- 15 was allowed to warm to room temperature. The product was extracted into ether and washed with 1N hydrochloric acid (50 ml). The organics were washed with brine (50 ml), dried over MgSO<sub>4</sub> and then concentrated *in vacuo* to a red oil. The oil was purified through a small plug of silica gel using 3:1-2:1 hexanes/ethyl acetate as eluent. Concentration of the column
- 20 fractions *in vacuo* gave 2-(3-iodobenzoyl)-acetonitrile (8.3 g) as a yellow oil.

**Step 2**

- A mixture of 2-(3-iodobenzoyl)acetonitrile (36.2 g, 133.5 mmol) and N,N-diphenylformamidine (26.2 g, 133.5 mmol) in toluene (200 ml) was heated at reflux under a nitrogen atmosphere. After 8 h, the reaction
- 25 mixture was cooled to room temperature and diluted with ether (200 ml) to give 2-(3-iodobenzoyl)-3-phenylaminoacrylonitrile (31.2 g) as a solid.

**Step 3**

A mixture of 4-fluorophenylhydrazine (26.6 g, 211 mmol) and 2-(3-iodobenzoyl)-3-phenylaminoacrylonitrile (79 g, 211 mmol) in ethanol (400

- 85 -

ml) was heated at reflux under a nitrogen atmosphere. After 30 minutes, the reaction mixture was cooled to room temperature, diluted with hexane to give 5-amino-4-(3-iodobenzoyl)-1-(4-fluorophenyl)pyrazole (75.1 g) as a solid.

5 Replacing 4-fluorophenylhydrazine with 4-methylphenylhydrazine, 3-methoxyphenylhydrazine, 4-sulfamoylphenylhydrazine, 2,4-dimethylphenylhydrazine, 2-methylphenylhydrazine, 4-chloro-2-methylphenylhydrazine, 4-methylsulfonylphenylhydrazine, 2-ethylphenylhydrazine, and 2,4-difluorophenylhydrazine in Step 3 above  
10 gave respectively:


5-amino-4-(3-iodobenzoyl)-1-(4-methylphenyl)pyrazole,  
5-amino-4-(3-iodobenzoyl)-1-(3-methoxyphenyl)pyrazole,  
5-amino-4-(3-iodobenzoyl)-1-(4-sulfamoylphenyl)pyrazole,  
5-amino-4-(3-iodobenzoyl)-1-(2,4-dimethylphenyl)pyrazole,  
15 5-amino-4-(3-iodobenzoyl)-1-(2-methylphenyl)pyrazole,  
5-amino-4-(3-iodobenzoyl)-1-(4-chloro-2-methylphenyl)pyrazole,  
5-amino-4-(3-iodobenzoyl)-1-(4-methylsulfonylphenyl)pyrazole,  
5-amino-4-(3-iodobenzoyl)-1-(2-ethylphenyl)pyrazol, and  
5-amino-4-(3-iodobenzoyl)-1-(2,4-difluorophenyl)pyrazole.

20

**Example 18**

5-amino-1-(4-fluorophenyl)-4-[3-(1,2-dihydroxyethyl)benzoyl] pyrazole (85)

5

**Step 1**

To a solution of 5-amino-1-(4-fluorophenyl)-4-[3-iodobenzoyl]pyrazole  
10 (10g, 24.6 mmol) in 100 ml dimethylformamide was added vinyltributyltin  
(8.57g, 27.0 mmol) and tetrakis(triphenylphosphine) palladium (0) (1.42g,  
1.23 mmol). The resulting solution was degassed with argon and  
subsequently warmed to 100°C for 12 hours.

The reaction was cooled to room temperature and was poured into  
15 500 ml distilled water and was extracted 3x 100 ml 1:1 ether/ethyl acetate.  
The organics were washed with brine (150 ml), dried over  $\text{MgSO}_4$  and then  
concentrated *in vacuo* to a brown oil. The oil was purified by flash column  
chromatography using 5:1-4:1 hexanes/ethyl acetate to remove impurities  
and 3:1-2:1 hexanes/ethyl acetate to elute the desired product.

20 Concentration of the column fractions *in vacuo* gave 5-amino-1-(4-  
fluorophenyl)-4-[3-vinylbenzoyl] pyrazole (4.48 g) as a white solid.

**Step 2**

To a suspension of 5-amino-1-(4-fluorophenyl)-4-[3-  
vinylbenzoyl] pyrazole (4.48g, 13.95 mmol) in 50 ml t-butanol was added  
25 N-methylmorpholine N-oxide (1.79g, 15.35 mmol) in 50 ml distilled water.  
To this mixture at room temperature was added a solution of 2.5% osmium  
tetraoxide in t-butanol (5.25 ml, 0.42 mmol). After 5 hours, the  
homogenous reaction was diluted with ethyl acetate (25 ml) and the  
organics were separated and washed with brine (25 ml), dried over  $\text{MgSO}_4$



- 87 -


and then concentrated *in vacuo* to a brown oil. The oil was purified by flash column chromatography using 1:1 hexanes/ethyl acetate to remove impurities and ethyl acetate to elute the desired product. Concentration of the column fractions *in vacuo* gave 5-amino-1-(4-fluorophenyl)-4-[3-(1,2-dihydroxyethyl)benzoyl] pyrazole (4.48 g) as a white foam. The foam was trituated to a solid from hexanes (2.36 g).

Replacing 5-amino-1-(4-fluorophenyl)-4-[3-iodobenzoyl]pyrazole in Step 1 above with:

5-amino-1-(2,4-difluorophenyl)-4-[3-iodobenzoyl]pyrazole and  
5-amino-1-(2-methylphenyl)-4-[3-iodobenzoyl]pyrazole, gave respectively  
5-amino-1-(2,4-difluorophenyl)-4-[3-(1,2-dihydroxyethyl)benzoyl] pyrazole (103) and  
5-amino-1-(2-methylphenyl)-4-[3-(1,2-dihydroxyethyl)benzoyl] pyrazole (109).

### Example 19

5-Amino-1-(2,4-difluorophenyl)-4-[3-(1-piperidinylmethyl)-benzoyl]pyrazole (86)



#### Step 1:

To a suspension of 5-amino-1-(2,4-difluorophenyl)-4-[3-(1,2-dihydroxyethane)benzoyl] pyrazole (10.1g, 28 mmol) in 100 ml t-butanol was added 100 ml distilled water and sodium periodate (18.06g, 84 mmol). After 2 hours, the solid precipitate was collected by vacuum filtration and was washed with 300 ml distilled water and dried *in vacuo* to give 8.28 g of

- 88 -

5-amino-1-(2,4-difluorophenyl)-4-[3-formylbenzoyl] pyrazole as a white solid.

Step 2:

To a solution of 5-amino-1-(2,4-difluorophenyl)-4-[3-formylbenzoyl] pyrazole (0.3 g, 0.92 mmol), piperidine (0.1 ml, 1.0 mmol), acetic acid (0.05 ml) in 1,2-dichloroethane (5 ml) was added sodium triacetoxyborohydride (0.29g, 1.37 mmol). After stirring at room temperature for 12 hours, the reaction was diluted with 10% hydrochloric acid and ethyl acetate (10 ml). The aqueous layer was separated and neutralized to pH 9 with sodium hydroxide and was then extracted with ethyl acetate. The combined organics were separated and washed with brine (25 ml), dried over  $\text{MgSO}_4$  and then concentrated *in vacuo* to a brown oil. The oil was purified by flash column chromatography using 1:1 hexanes/ethyl acetate to remove impurities and ethyl acetate to elute the desired product. Concentration of the column fractions *in vacuo* gave 5-amino-1-(4-fluorophenyl)-4-[3-(1-piperidinylmethyl)-benzoyl] pyrazole as an oil (0.211g). The compound was triturated to a solid from hexanes/ethyl acetate.

Replacing piperidine in Step1 above with:

morpholine,

N-methylpiperazine,

4-hydroxypiperidine,

2-aminopyridine,






3-aminopyridine,

4-methylimidazole,




3-aminopyrazole, and

2-methylimidazole;

the following compounds were obtained


Mole structure	CPD#	HRMS MWt	M. Pt.
	131	398.411	127.3-128.5
 ClH ClH	132	484.376	238.2-238.6
	133	412.438	141.5-145.5
	134	405.406	
	135	405.406	

- 90 -

	136	393.395	
	137	508.405	
	138	507.417	

**Example 20****5-amino-1-(3-methylphenyl)-4-[3-{N-oxidopyridin-3-yl}benzoyl]pyrazole  
(70)**

5



A mixture of 5-amino-4-(3-iodobenzoyl)-1-(2-methylphenyl)pyrazole, from Example 17 (0.98 g, 2.4 mmol), pinacol diboron (0.68 g, 2.7 mmol),  
 10 [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (0.2 g, .24mmol) and potassium acetate (0.72 g, 7.3mmol) in DMF (10 ml) was heated at 80 degrees, under argon. After 2 h, the reaction mixture was cooled to room temperature and 3-bromopyridine N-oxide (.47 g, 2.7 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (0.2 g, .24 mmol) and 2  
 15 M sodium carbonate (6.1 ml, 12.2 mmol) was added and heated to 80 degrees. After 16 h, the reaction mixture was cooled to room temperature, poured into brine and the product extracted into ethyl acetate. The organic layer was dried over sodium sulfate, filtered and then the solution was evaporated to dryness. The residue was purified by flash chromatography (gradient elution: ethyl acetate to 20 % methanol/ethyl acetate) to give,  
 20 after recrystallization from methanol/ethyl acetate/hexane, 5-amino-1-(3-methylphenyl)-4-[3-(N-oxidopyridin-3-yl)benzoyl]pyrazole (0.57 g, mpt. 190.5-191.2).

Replacing 5-amino-4-(3-iodobenzoyl)-1-(2-methylphenyl)pyrazole/3-bromopyridine N-oxide with:  
 25 5-amino-4-(3-iodobenzoyl)-1-(4-methylphenyl)pyrazole/3-bromopyridine,  
 5-amino-4-(3-iodobenzoyl)-1-(3-methoxyphenyl)pyrazole/3-bromopyridine,  
 5-amino-4-(3-iodobenzoyl)-1-(4-sulfamoylphenyl)pyrazole/3-bromopyridine,

- 92 -

5-amino-4-(3-iodobenzoyl)-1-(2,4-dimethylphenyl)pyrazole/3-bromopyridine,

5-amino-4-(3-iodobenzoyl)-1-(2-methylphenyl)pyrazole/3-bromopyridine-N-oxide,

5 5-amino-4-(3-iodobenzoyl)-1-(4-chloro-2-methylphenyl)pyrazole/3-bromopyridine,

5-amino-4-(3-iodobenzoyl)-1-(4-methylsulfonylphenyl)pyrazole/3-bromopyridine,

5-amino-4-(3-iodobenzoyl)-1-(2-ethylphenyl)pyrazole/3-bromopyridine, and

10 5-amino-4-(3-iodobenzoyl)-1-(2,4-difluorophenyl)pyrazole/2-bromoimidazole,

gave respectively the following compounds (as their hydrochloride salts as appropriate):

15

5-amino-1-(4-methylphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (65),

5-amino-1-(3-methoxyphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (66),

5-amino-1-(4-sulfamoylphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (68),

5-amino-1-(2,4-dimethylphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (69),

20 5-amino-1-(2-methylphenyl)-4-[3-(N-oxidopyridin-3-yl)benzoyl]pyrazole (70),

5-amino-1-(4-chloro-2-methylphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (73),

25 5-amino-1-(4-methylsulfonylphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (75),

5-amino-1-(2-ethylphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (76), and


5-amino-1-(2,4-difluorophenyl)-4-[3-(imidazol-2-yl)benzoyl]pyrazole (77).

### Example 21

30

5-amino-1-(2,4-difluorophenyl)-4-[N-oxidopyridin-3-yl]benzoyl]pyrazole (60)

- 93 -




To a solution of 5-amino-1-(2,4-difluorophenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole (4.6 g, 12.2 mmol) in dichloromethane (100ml) was  
5 added 3-chloroperoxybenzoic acid (5.6 g, 18.3 mmol) and the mixture was stirred at room temperature. After 4 h, a solution of 10 % aqueous sodium sulfite (50 ml) was added. After 0.5 h, the organic layer was separated, washed with brine, dried over sodium sulfate and filtered. The filtrate was  
10 concentrated to dryness and the residue purified by flash chromatography (gradient elution: ethyl acetate to 30 % methanol/ethyl acetate) to give, after recrystallization from methanol, 5-amino-1-(2,4-difluorophenyl)-4-[3-(N-oxidopyridin-3-yl) benzoyl]pyrazole (1.3g, Mpt. 251.1-251.7°C).

- 94 -

**Example 22**

5-amino-1-(2,4-difluorophenyl)-4-[3-(pyridin-4-yl)benzoyl]pyrazole (61)

5




A mixture of 5-amino-4-(3-bromobenzoyl)-1-(2,4-difluorophenyl)pyrazole (.93 g, 2.5 mmol), 4-tributylstannylpyridine (1.0 g, 2.7 mmol) and bis(triphenylphosphine)palladium chloride (.17 g, 2.5 mmol) in DMF (15ml) was heated at 100 degrees under argon. After 16 h, the reaction mixture was cooled to room temperature and a solution of 10 % aqueous potassium fluoride (30 ml) was added. After 1 h, the reaction mixture was poured into brine, extracted with ethyl acetate, dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by flash chromatography (gradient elution: 50-100 % ethyl acetate/hexane to 5 % methanol/ethyl acetate to give, after recrystallizaion from methanol/ethyl acetate, 5-amino-1-(2,4-difluorophenyl)-4-[3-(pyridin-4-yl)benzoyl]pyrazole (0.42 g, Mpt. 218-226 °C).



**Example 23**

5-Amino-1-(2,4-dimethylphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole HCl  
salt (69)

5

**Step 1**

10        Into a solution of n-butyl lithium (165 ml, 264 mmol) in butyl ether  
(250 ml) at -78 degrees under nitrogen was added 3-bromopyridine (25.4  
ml, 264 mmol). After 1 h, added diethylmethoxyborane (52ml, 396 mmol).  
The mixture was allowed to warm to room temperature. After 16 h, added  
water and brine, separated organic layer, dried over sodium sulfate, then  
15        concentrated. The resulting slurry was dissolved in isopropanol (500 ml),  
cooled and the product isolated by filtration give diethyl(3-pyridyl)borane  
(29.8g).

**Step 2**

20        A mixture of diethyl(3-pyridyl)borane (176.4g, 1.2mole), methyl-3-  
iodobenzoate(262g, 1 mole), potassium phosphate (318.4 g, 1.5 mole) and  
tetrakis(triphenyl)phosphine palladium (0) (57.8g, .05 mole) in DMF (1000  
ml) was heated at 80 degrees under argon. After 10 h, the mixture was  
diluted with water and extracted with ethyl acetate. The organic layer was  
filtered and washed with water. To the organic fraction was added  
25        concentrated HCL (65 ml) . The organic layer was separated and extracted  
with aqueous HCl. The combined acid extractions were treated with ethyl  
acetate, followed by 50% aqueous sodium hydroxide (55 ml). The organic

- 96 -

layer was separated, washed with water and saturated sodium bicarbonate solution, then dried over sodium sulfate. The solution was filtered and concentrated to give methyl-3-(pyridin-3-yl)benzoate (145.3g).

### Step 3

5 To a solution of methyl-3-(pyridin-3-yl)benzoate (126.2 g, 0.59 mole) in THF (600 ml) was added acetonitrile (37 ml, 0.71 mole) and the reaction was cooled to - 40 degrees. A solution of lithium diisopropyl amide (590 ml, 1.18 mole) was added dropwise. After 30 minutes, added methanol (25 ml) and after another 30 minutes, added water (110 ml). Allowed the reaction  
10 mixture to warm to 10 degrees and added ethyl acetate. The layers were separated and the aqueous layer was acidified with 1 M HCl. The aqueous layer was extracted with ethyl acetate, diluted with hexane and washed with brine. The organic phase was concentrated, then combined with N,N'-diphenylformanidine (120 g, 0.61 mole) in 800 ml of ethyl acetate. The  
15 mixture was stirred at room temperature. After 3 days, the product was collected by filtration and recrystallized from isopropanol, hexane to give 2-(3-pyridin-3-yl)phenyl-3-phenylacrylonitrile (100 g).




### Step 4

A solution of 2-(3-pyridin-3-ylphenyl)-3-phenylacrylonitrile (1.0 g, 3  
20 mmol) and 2,4-dimethylphenylhydrazine (0.4 g, 3 mmol) in ethanol (30 ml) was heated at reflux, under nitrogen. After 6 h, the reaction was cooled to room temperature, concentrated to dryness and the residue purified by flash column chromatography (elution gradient:40-100% ethyl  
acetate/hexane to 10% methanol/ethyl acetate). The purified residue was  
25 taken up in ethyl acetate and HCl/ether added to prepare the salt. After recrystallization from methanol/ethyl acetate was isolated 5-amino-1-(2,4-dimethylphenyl)-4-[3-(pyridin-3-yl)benzoyl]pyrazole hydrochloride salt (0.74 g, m.pt. 250.7-251.8).




Proceeding as above in Example 23, but replacing 2,4-  
30 dimethylphenylhydrazine in step 4 with:  
phenylhydrazine,  
2-methyl-4-chlorophenylhydrazine,  
4-methoxyphenylhydrazine,  
4-methylsulfonylphenylhydrazine,





- 98 -

 ClH	140	406.871
 ClH	75	454.936
 ClH	76	404.899

- 99 -

 <p>ClH</p>	141	418.926
 <p>ClH</p>	142	406.871
 <p>ClH</p>	143	425.317


- 100 -

 ClH	144	394.835
 ClH	145	408.862

**Example 24**

5-amino-1-(4-fluorophenyl)-4-[3-{2(R),3-dihydroxypropoxy}benzoyl] pyrazole  
(106)

5

**Step 1**

To a solution of 5-amino-4-(3-hydroxybenzoyl)-1-(4-fluorophenyl)pyrazole (0.5g, 1.68 mmol) in 5 ml dry dimethylformamide was added D- $\alpha$ ,  $\beta$ -isopropylideneglycerol- $\gamma$ -tosylate (0.72g, 2.52 mmol) followed by anhydrous potassium carbonate (0.695g, 5.04 mmol). The reaction was warmed to 80 °C under argon. After 24 hours, the reaction was cooled to room temperature and an additional 500 mg of D- $\alpha$ , $\beta$ -isopropylideneglycerol- $\gamma$ -tosylate was added and the reaction was warmed back to 80 °C under argon. After 8 additional hours, the reaction was cooled to room temperature and diluted with distilled water (50 ml) and the product was extracted into ether. The combined organics were washed with brine (50 ml), dried over MgSO<sub>4</sub> and then concentrated *in vacuo* to a yellow oil. The oil was purified by flash column chromatography on silica gel using 2:1-1:1 hexanes/ethyl acetate as eluent. Concentration of the column fractions *in vacuo* gave 556 mg of the desired acetal.

**Step 2**


To a solution of the acetal formed above (0.556 g, 1.35 mmol) in methanol (10 ml) was added distilled water (2 ml) and p-toluenesulfonic acid monohydrate (5 mg). The solution was warmed to 80 °C under an

- 102 -

argon atmosphere. After 2 h, the reaction mixture was cooled to room temperature concentrated *in vacuo* to a yellow oil which was redissolved in ethyl acetate (50 ml) and saturated sodium bicarbonate (50 ml). The organic layer was separated, dried over  $\text{MgSO}_4$  and then concentrated *in vacuo* to a white solid. Recrystallization from hexanes/ethyl acetate gave 196 mg of the desired diol (Mpt. 150.2-153.0°C).

### Example 25

5-amino-1-(4-fluorophenyl)-4-thenoyl-pyrazole (114)



Proceeding as described in Example 1, step 2, but replacing 2-(3-bromobenzoyl)acetonitrile with 2-(2-thenoyl)acetonitrile and then following step 3 was obtained 5-amino-1-(4-fluorophenyl)-4-(2-thenoyl)-pyrazole.

Proceeding as described in Example 1, step 2, but replacing 2-(3-bromobenzoyl)acetonitrile with 2-(2-furanoyl)acetonitrile and then following step 3 was obtained 5-amino-1-(4-fluorophenyl)-4-(2-furanoyl)-pyrazole (115).

Proceeding as described in Example 1, step 2, but replacing 2-(3-bromobenzoyl)acetonitrile with 2-(2-methyl-3-furanoyl)acetonitrile and substituting 4-fluorophenylhydrazine in step 3 with 2,4-difluorophenylhydrazine was obtained 5-amino-1-(2,4-difluorophenyl)-4-(2-methylfuran-3-oyl)-pyrazole (116).

Proceeding as described in Example 1, step 2, but replacing 2-(3-bromobenzoyl)acetonitrile with 2-(6-quinolinoyl)acetonitrile and substituting 4-fluorophenylhydrazine in step 3 with 2,4-difluorophenylhydrazine was obtained 5-amino-1-(2,4-difluorophenyl)-4-(6-quinolinoyl)-pyrazole.HCl (117) (mpt. 220-259.2).



**Example 26**

The following are representative pharmaceutical formulations containing a compound of Formula (I).

5

**Tablet formulation**

The following ingredients are mixed intimately and pressed into single scored tablets.

10

Ingredient	Quantity per tablet, mg
compound of this invention	400
cornstarch	50
croscarmellose sodium	25
lactose	120
magnesium stearate	5

15

**Capsule formulation**

20

The following ingredients are mixed intimately and loaded into a hard-shell gelatin capsule.

Ingredient	Quantity per capsule, mg
compound of this invention	200
lactose, spray-dried	148
magnesium stearate	2

25

- 104 -

Suspension formulation

The following ingredients are mixed to form a suspension for oral administration.

5

Ingredient	Amount
compound of this invention	1.0 g
fumaric acid	0.5 g
sodium chloride	2.0 g
methyl paraben	0.15 g
propyl paraben	0.05 g
granulated sugar	25.5 g
sorbitol (70% solution)	12.85 g
Veegum K (Vanderbilt Co.)	1.0 g
flavoring	0.035 ml
colorings	0.5 mg
distilled water	q.s. to 100 ml

15

Injectable formulation

20

The following ingredients are mixed to form an injectable formulation.

25

Ingredient	Amount
compound of this invention	0.2 g
sodium acetate buffer solution, 0.4 M	2.0 ml
HCl (1N) or NaOH (1N)	q.s. to suitable pH
water (distilled, sterile)	q.s. to 20 ml

30

All of the above ingredients, except water, are combined and heated to 60-70 °C with stirring. A sufficient quantity of water at 60 °C is then added with vigorous stirring to emulsify the ingredients, and water then added q.s. to 100 g.

- 105 -

Suppository formulation

A suppository of total weight 2.5 g is prepared by mixing the compound of the invention with Witepsol® H-15 (triglycerides of saturated vegetable fatty acid; Riches-Nelson, Inc., New York), and has the following composition:

compound of the invention	500 mg
Witepsol® H-15	balance

**Example 27**Inhibition Of p-38 (MAP) Kinase...*In Vitro* Assay

The p-38 MAP kinase inhibitory activity of compounds of this invention *in vitro* was determined by measuring the transfer of the  $\gamma$ -phosphate from  $\gamma$ -<sup>33</sup>P-ATP by p-38 kinase to Myelin Basic Protein (MBP), using the a minor modification of the method described in Ahn, N. G.; et al. *J. Biol. Chem.* Vol. **266**(7), 4220-4227, (1991).

The phosphorylated form of the recombinant p38 MAP kinase was expressed with SEK-1 and MEKK in E. Coli and then purified by affinity chromatography using a Nickel column.

The phosphorylated p38 MAP kinase was diluted in kinase buffer (20 mM 3-(N-morpholino)propanesulfonic acid, pH 7.2, 25 mM  $\beta$ -glycerol phosphate, 5 mM ethylene glycol-bis(beta-aminoethyl ether)-N,N,N',N'-tetraacetic acid, 1mM sodium vanadate, 1mM dithiothreitol, 40mM magnesium chloride). Test compound dissolved in DMSO or only DMSO (control) was added and the samples were incubated for 10 min at 30 °C. The kinase reaction was initiated by the addition of a substrate cocktail containing MBP and  $\gamma$ -<sup>33</sup>P-ATP. After incubating for an additional 20 min at 30 °C, the reaction was terminated by adding 0.75% phosphoric acid. The phosphorylated MBP was then separated from the residual  $\gamma$ -<sup>33</sup>P-ATP using a phosphocellulose membrane (Millipore, Bedford, MA) and quantitated using a scintillation counter (Packard, Meriden, CT).

Compounds of the invention wer active in this assay. The p-38 inhibitory activities (expressed as IC<sub>50</sub> , the concentration causing 50 %

- 106 -

inhibition of the p-38 enzyme being assayed) of some compounds of the invention are:

CPD #	IC <sub>50</sub> , $\mu$ M	CPD #	IC <sub>50</sub> , $\mu$ M
1	1.81	19	1.45
2	3.29	21	2.18
3	1.78	27	2.72
4	6.18	33	1.12
6	1.74	38	6.31
9	1.32	43	6.52
14	1.27	50	1.25

5

### Example 28

#### Inhibition of LPS-Induced TNF- $\alpha$ Production In THP1 Cells....*In Vitro* Assay

10 The ability of the compounds of this invention to inhibit the TNF- $\alpha$  release was determined using a minor modification of the methods described in Blifeld, C. et al. *Transplantation*, Vol. 51(2), 498-503, (1991).

##### (a) Induction of TNF biosynthesis:

15 THP-1 cells were suspended in culture medium [RPMI (Gibco-BRL, Gaithersburg, MD) containing 15% fetal bovine serum, 0.02 mM 2-mercaptoethanol], at a concentration of  $2.5 \times 10^6$  cells/ml and then plated in 96 well plate (0.2 ml aliquots in each well). Test compounds were dissolved in DMSO and then diluted with the culture medium such that the final DMSO concentration was 5%. 20  $\mu$ l aliquots of test solution or only medium with DMSO (control) were added to each well. The cells were incubated for 30 min., at 37 °C. LPS (Sigma, St. Louis, MO) was added to the wells at a final concentration of 0.5  $\mu$ g/ml, and cells were incubated for an additional 2 h. At the end of the incubation period, culture supernatants were collected and the amount of TNF- $\alpha$  present was determined using an ELISA assay as described below.

##### 25 (b) ELISA Assay:

The amount of human TNF- $\alpha$  present was determined by a specific trapping ELISA assay using two anti-TNF- $\alpha$  antibodies (2TNF-H22 and 2TNF-H34) described in Reimund, J. M., et al. *GUT*. Vol. 39(5), 684-689 (1996).

5 Polystyrene 96-well plates were coated with 50  $\mu$ l per well of antibody 2TNF-H22 in PBS (10  $\mu$ g/ml) and incubated in a humidified chamber at 4 °C overnight. The plates were washed with PBS and then blocked with 5% nonfat-dry milk in PBS for 1 hour at room temperature and washed with 0.1% BSA (bovine serum albumin) in PBS.

10 TNF standards were prepared from a stock solution of human recombinant TNF- $\alpha$  (R&D Systems, Minneapolis, MN). The concentration of the standards in the assay began at 10 ng/ml followed by 6 half log serial dilution's.

25  $\mu$ l aliquots of the above culture supernatants or TNF standards  
15 or only medium (control) were mixed with 25  $\mu$ l aliquots of biotinylated monoclonal antibody 2TNF-H34 (2  $\mu$ g/ml in PBS containing 0.1% BSA) and then added to each well. The samples were incubated for 2 h at room temperature with gentle shaking and then washed 3 times with 0.1% BSA in PBS. 50  $\mu$ l of peroxidase-streptavidin (Zymed, S. San Francisco, CA) solution containing 0.416  $\mu$ g/ml  
20 of peroxidase-streptavidin and 0.1% BSA in PBS was added to each well. The samples were incubated for an additional 1 h at room temperature and then washed 4 times with 0.1% BSA in PBS. 50  $\mu$ l of O-phenylenediamine solution (1 $\mu$ g/ml O-phenylene-diamine and 0.03 % hydrogen peroxide in  
25 0.2M citrate buffer pH 4.5) was added to each well and the samples were incubated in the dark for 30 min., at room temperature. Optical density of the sample and the reference were read at 450 nm and 650 nm, respectively. TNF- $\alpha$  levels were determined from a graph relating the optical density at 450 nm to the concentration used.

30 The IC<sub>50</sub> value was defined as the concentration of the test compound corresponding to half-maximal reduction in 450 nm absorbance. Compounds of the invention were active in this assay. The activity of selected compounds is shown below.

- 108 -

CPD #	IC <sub>50</sub> , $\mu$ M	CPD #	IC <sub>50</sub> , $\mu$ M
1	1.77	21	0.61
2	6.30	27	0.83
4	1.26	33	0.14
6	1.04	38	0.69
10	1.62	43	0.17
13	0.77	50	0.51
19	0.17		

### Example 29

#### Inhibition of LPS-Induced TNF- $\alpha$ Production In Rats....*In Vivo* Assay

5        The ability of the compounds of this invention to inhibit the TNF- $\alpha$  release, *in vivo*, was determined using a minor modification of the methods described in described in Zanetti, G.; Heumann, D., *et. al.*, "Cytokine production after intravenous or peritoneal Gram-negative bacterial challenge in mice," *J. Immunol.*, **148**, 1890, (1992) and Sekut, L., Menius, J.A., *et. al.*, "Evaluation of the significance of elevated levels of systemic and localized tumor necrosis factor in different animal models of inflammation," *J. Lab. Clin. Med.*, **124**, 813, (1994).

15        Female Sprague-Dawley rats weighing 110-140 grams (Charles River, Hollister, CA) were acclimated for one week. Groups containing 8 mice each were dosed orally either with the test compounds dissolved in an aqueous vehicle containing 0.9% sodium chloride, 0.5% sodium carboxymethyl-cellulose, 0.4% polysorbate 80, 0.9% benzyl alcohol (CMC vehicle) or only vehicle (control group). After 30 min., the mice were injected intraperitoneally with 50  $\mu$ g/kg of LPS (Sigma, St. Louis, MO).

20        After 1.5 h, the mice were sacrificed by CO<sub>2</sub> inhalation and blood was harvested by cardiocentesis. Blood was clarified by centrifugation at 15,600 X g for 5 min., and sera were transferred to clean tubes and frozen at -20°C until analyzed for TNF- $\alpha$  by ELISA assay (Biosource International, Camarillo, CA) following the manufacturer's protocol.

- 109 -

The TNF- $\alpha$  inhibitory activity of selected compounds of the invention, i.e., the measure of the TNF- $\alpha$  content in the test group relative to the vehicle treated group (control group) at 30 mg was:

CPD #	% Inhibiti on	CPD #	% Inhibiti on
3	96	19	76
8	86	34	75
16	86		

5

### Example 30

#### Adjuvant Arthritis Assay In Rats.....In Vivo assay

The Anti-inflammatory activity of the compounds of this invention was determined utilizing adjuvant induced arthritis in rats. Briefly, Female Sprague Dawley rats, weighing 120-155 g (Charles River, Hollister, CA) were acclimated in-house for approximately 1 week prior to use. On day 1, the animals were injected intradermally in the 1/4 proximal portion of the tail with 0.1 ml of a mineral oil (Sigma, St. Louis, MO) suspension of heat killed and dried *Mycobacterium Butyricum* (Difco, Bacto., Des., Lot 115979JA/EXP9/99) at a concentration of 1mg/0.1ml.

On day 7, the test compounds were administered in CMC vehicle through to day 18. On day 18, following the administration of the compound, animals were weighed. Clinical scores were obtained to evaluate the intensity of edema in the four paws and tail. A score of 0 to 4 was assigned to each paw and 0 to 3 to the tail such that the maximum score was 19. Polyarthritic animals were scored 0 when no inflammatory signs ( swelling and redness ) were observed in any of the small joints (intrapthalangeal, metacarpophalangeal, metatarsophalangeal) or large joints (wrist/ carpus, ankle/tarsus). Animals were scored 1 when slight inflammation was observed, 2 moderate edema, 3 severe edema, and 4 when very severe edema was present. The tail was scored 0 when no signs of edema or necrotic tissue was observed, 1 when inocula injection sites

- 110 -

and immediate surrounding tissue exhibit slight edema, 2 when approximately 1/4 of the tail was either inflamed or exhibiting necrotic tissue, and 3 when over 1/4 of the tail exhibited severe necroses or edema. Following clinical scores, the hind paws were transected at the distal tibia, 5 just proximal to the tarsal joint. The left and right hind paws were weighed individually, and recorded.

The compounds of the present invention exhibit anti-inflammatory activity when tested in this assay.


10 The foregoing invention has been described in some detail by way of illustration and example, for purposes of clarity and understanding. It will be obvious to one of skill in the art that changes and modifications may be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive.



- 111 -

Claims

1. A compound selected from the group of compounds represented by Formula (I):



wherein:

$R^1$  is hydrogen, acyl or  $-P(O)(OH)_2$ ;

$R^2$  is hydrogen, halo, alkyl or alkylthio;

A is an aryl, heteroaryl or a heterocyclyl ring optionally fused to a phenyl ring provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom;

B is an aryl or heteroaryl ring;

$R^3$  is selected from the group consisting of:

- (a) amino, alkylamino or dialkylamino;
- (b) acylamino;
- (c) optionally substituted heterocyclyl;
- (d) optionally substituted aryl or heteroaryl;
- (e) heteroalkyl;
- (f) heteroalkenyl;
- (g) heteroalkynyl;
- (h) heteroalkoxy;
- (i) heteroalkylamino;
- (j) optionally substituted heterocyclylalkyl;
- (k) optionally substituted heterocyclylalkenyl;
- (l) optionally substituted heterocyclylalkynyl;
- (m) optionally substituted heterocyclylalkoxy or heterocycliloxy;

- 112 -

- (n) optionally substituted heterocyclalkylamino;
- (o) optionally substituted heterocyclalkylcarbonyl;
- (p) heteroalkylcarbonyl;
- (q)  $-\text{NHSO}_2\text{R}^6$  where  $\text{R}^6$  is alkyl, heteroalkyl or  
 5 optionally substituted heterocyclalkyl;
- (r)  $-\text{NHSO}_2\text{NR}^7\text{R}^8$  where  $\text{R}^7$  and  $\text{R}^8$  are,  
 independently of each other, hydrogen, alkyl or  
 heteroalkyl;
- (s)  $-\text{Y}-(\text{alkylene})-\text{R}^9$  where:  
 10 Y is a single bond,  $-\text{O}-$ ,  $-\text{NH}-$  or  $-\text{S}(\text{O})_n-$  (where n is  
 an integer from 0 to 2); and  
 $\text{R}^9$  is cyano, optionally substituted heteroaryl,  $-\text{COOH}$ ,  $-\text{COR}^{10}$ ,  $-\text{COOR}^{11}$ ,  $-\text{CONR}^{12}\text{R}^{13}$ ,  $-\text{SO}_2\text{R}^{14}$ ,  $-\text{SO}_2\text{NR}^{15}\text{R}^{16}$ ,  $-\text{NHSO}_2\text{R}^{17}$  or  $-\text{NHSO}_2\text{NR}^{18}\text{R}^{19}$ , where  
 15  $\text{R}^{10}$  is alkyl or optionally substituted heterocycle,  
 $\text{R}^{11}$  is alkyl, and  $\text{R}^{12}$ ,  $\text{R}^{13}$ ,  $\text{R}^{14}$ ,  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{17}$ ,  $\text{R}^{18}$  and  
 $\text{R}^{19}$  are, independently of each other, hydrogen,  
 alkyl or heteroalkyl;
- (t)  $-\text{C}(=\text{NR}^{20})(\text{NR}^{21}\text{R}^{22})$  where  $\text{R}^{20}$ ,  $\text{R}^{21}$  and  $\text{R}^{22}$   
 20 independently represent hydrogen, alkyl or  
 hydroxy, or  $\text{R}^{20}$  and  $\text{R}^{21}$  together are  $-(\text{CH}_2)_n-$   
 where n is 2 or 3 and  $\text{R}^{22}$  is hydrogen or alkyl;
- (u)  $-\text{NHC}(\text{X})\text{NR}^{23}\text{R}^{24}$  where X is  $-\text{O}-$  or  $-\text{S}-$ , and  $\text{R}^{23}$   
 25 and  $\text{R}^{24}$  are, independently of each other,  
 hydrogen, alkyl or heteroalkyl;
- (v)  $-\text{CONR}^{25}\text{R}^{26}$  where  $\text{R}^{25}$  and  $\text{R}^{26}$  independently  
 represent hydrogen, alkyl, heteroalkyl or  
 optionally substituted heterocyclalkyl, or  $\text{R}^{25}$   
 and  $\text{R}^{26}$  together with the nitrogen to which they  
 30 are attached form an optionally substituted  
 heterocycl ring;
- (w)  $-\text{S}(\text{O})_n\text{R}^{27}$  where n is an integer from 0 to 2, and  
 $\text{R}^{27}$  is alkyl, heteroalkyl, optionally substituted  
 heterocyclalkyl or

- 113 -

-NR<sup>28</sup>R<sup>29</sup> where R<sup>28</sup> and R<sup>29</sup> are, independently of each other, hydrogen, alkyl or heteroalkyl;

- (x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;
- (y) arylaminoalkylene or heteroarylaminomethylene;
- (z) Z-alkylene-NR<sup>30</sup>R<sup>31</sup> or Z-alkylene-OR<sup>32</sup> where Z is -NH-, -N(lower alkyl)- or -O-, and R<sup>30</sup>, R<sup>31</sup> and R<sup>32</sup> are independently of each other, hydrogen, alkyl or heteroalkyl;
- (aa) -OC(O)-alkylene-CO<sub>2</sub>H or -OC(O)-NR'R'' (where R' and R'' are independently hydrogen or alkyl); and
- (bb) heteroarylalkenylene or heteroarylalkynylene;

R<sup>4</sup> is selected from the group consisting of:

- (a) hydrogen;
- (b) halo;
- (c) alkyl;
- (d) alkoxy; and
- (e) hydroxy;

R<sup>5</sup> is selected from the group consisting of :

- (a) hydrogen;
- (b) halo;
- (c) alkyl;
- (d) haloalkyl;
- (e) thioalkyl;
- (f) hydroxy;
- (g) amino;
- (h) alkylamino;
- (i) dialkylamino;
- (j) heteroalkyl;
- (k) optionally substituted heterocycle;
- (l) optionally substituted heterocyclalkyl;
- (m) optionally substituted heterocyclalkoxy;

- 114 -

- (n) alkylsulfonyl;
- (o) aminosulfonyl, mono-alkylaminosulfonyl or di-alkylaminosulfonyl;
- (p) heteroalkoxy; and
- (q) carboxy;

R<sup>6</sup> is selected from the group consisting of:

- (a) hydrogen;
- (b) halo;
- (c) alkyl; and
- (d) alkoxy;

prodrugs, individual isomers, mixtures of isomers and pharmaceutically acceptable salts thereof.

2. A compound according to claim 1, wherein:

- R<sup>1</sup> is hydrogen or acyl;
- R<sup>2</sup> is hydrogen or alkyl;
- A is an aryl or heteroaryl ring;

3. A compound according to claim 1, wherein:

- R<sup>1</sup> is hydrogen, acyl or -P(O)(OH)<sub>2</sub>;
- R<sup>2</sup> is hydrogen, halo, alkyl or alkylthio;
- A is an aryl, heteroaryl or a heterocyclyl ring optionally fused to a phenyl ring provided that the heterocyclyl ring is attached to the carbonyl group via a carbon ring atom;
- B is an aryl or heteroaryl ring;
- R<sup>3</sup> is selected from the group consisting of:

- (a) amino;
- (b) acylamino;
- (c) optionally substituted heterocycle;
- (d) heteroaryl optionally substituted with a substituent selected from halo, alkyl or alkoxy;
- (e) heteroalkyl;
- (f) heteroalkenyl;
- (g) heteroalkynyl;

- 115 -

- (h) heteroalkoxy;
- (i) heteroalkylamino;
- (j) optionally substituted heterocyclalkyl;
- (k) optionally substituted heterocyclalkenyl;
- (l) optionally substituted heterocyclalkynyl;
- (m) optionally substituted heterocyclalkoxy;
- (n) optionally substituted heterocyclalkylamino;
- (o) optionally substituted heterocyclalkylcarbonyl;
- (p) heteroalkylcarbonyl;
- (q)  $-\text{NHSO}_2\text{R}^6$  where  $\text{R}^6$  is alkyl, heteroalkyl or optionally substituted heterocyclalkyl;
- (r)  $-\text{NHSO}_2\text{NR}^7\text{R}^8$  where  $\text{R}^7$  and  $\text{R}^8$  are, independently of each other, hydrogen, alkyl or heteroalkyl;
- (s)  $-\text{Y}-(\text{alkylene})-\text{R}^9$  where:  
 $\text{Y}$  is a single bond,  $-\text{O}-$ ,  $-\text{NH}-$  or  $-\text{S}(\text{O})_n-$  (where  $n$  is an integer from 0 to 2); and  
 $\text{R}^9$  is cyano, heteroaryl,  $-\text{COOH}$ ,  $-\text{COR}^{10}$ ,  $-\text{COOR}^{11}$ ,  $-\text{CONR}^{12}\text{R}^{13}$ ,  $-\text{SO}_2\text{R}^{14}$ ,  $-\text{SO}_2\text{NR}^{15}\text{R}^{16}$ ,  $-\text{NHSO}_2\text{R}^{17}$  or  $-\text{NHSO}_2\text{NR}^{18}\text{R}^{19}$  where  $\text{R}^{10}$  is alkyl or optionally substituted heterocycle,  $\text{R}^{11}$  is alkyl, and  $\text{R}^{12}$ ,  $\text{R}^{13}$ ,  $\text{R}^{14}$ ,  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{17}$ ,  $\text{R}^{18}$  and  $\text{R}^{19}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;
- (t)  $-\text{C}(=\text{NR}^{20})(\text{NR}^{21}\text{R}^{22})$  where  $\text{R}^{20}$ ,  $\text{R}^{21}$  and  $\text{R}^{22}$  independently represent hydrogen, alkyl or hydroxy, or  $\text{R}^{20}$  and  $\text{R}^{21}$  together are  $-(\text{CH}_2)_n-$  where  $n$  is 2 or 3 and  $\text{R}^{22}$  is hydrogen or alkyl;
- (u)  $-\text{NHC}(\text{X})\text{NR}^{23}\text{R}^{24}$  where  $\text{X}$  is  $-\text{O}-$  or  $-\text{S}-$ , and  $\text{R}^{23}$  and  $\text{R}^{24}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;
- (v)  $-\text{CONR}^{25}\text{R}^{26}$  where  $\text{R}^{25}$  and  $\text{R}^{26}$  independently represent hydrogen, alkyl, heteroalkyl or optionally substituted heterocyclalkyl, or  $\text{R}^{25}$  and  $\text{R}^{26}$  together with the nitrogen to which they

- 116 -

are attached form an optionally substituted heterocyclyl ring;

- (w)  $-S(O)_nR^{27}$  where  $n$  is an integer from 0 to 2, and  $R^{27}$  is alkyl, heteroalkyl, optionally substituted heterocyclylalkyl or  $-NR^{28}R^{29}$  where  $R^{28}$  and  $R^{29}$  are, independently of each other, hydrogen, alkyl or heteroalkyl;

$R^4$  is selected from the group consisting of:

- (a) hydrogen;
- (b) halo;
- (c) alkyl; and
- (d) alkoxy;

$R^5$  is selected from the group consisting of :

- (a) hydrogen;
- (b) halo;
- (c) alkyl;
- (d) haloalkyl;
- (e) thioalkyl;
- (f) hydroxy;
- (g) amino;
- (h) alkylamino;
- (i) dialkylamino;
- (j) heteroalkyl;
- (k) optionally substituted heterocycle;
- (l) optionally substituted heterocyclylalkyl; and
- (m) optionally substituted heterocyclylalkoxy;

$R^6$  is selected from a group consisting of:

- (a) hydrogen;
- (b) halo;
- (c) alkyl; and
- (d) alkoxy

4. The compound of claim 1 or claim 2, wherein  $R^3$  is:

- (a) optionally substituted heterocyclyl;

- 117 -

- 5 (b) aryl or heteroaryl both optionally substituted with a substituent selected from halo, alkyl, amino, alkoxy, carboxy, lower alkoxy carbonyl,  $\text{SO}_2\text{R}'$  (where  $\text{R}'$  is alkyl) or  $\text{SO}_2\text{NHR}'\text{R}''$  (where  $\text{R}'$  and  $\text{R}''$  are independently hydrogen or alkyl);
- (c) heteroalkyl;
- (d) heteroalkenyl;
- (e) heteroalkylamino;
- (f) heteroalkoxy;
- 10 (g) optionally substituted heterocyclalkyl or heterocyclalkoxy;
- (h) optionally substituted heterocyclalkenyl;
- (i) optionally substituted heterocyclalkynyl;
- (j) optionally substituted heterocyclalkoxy;
- 15 (k) optionally substituted heterocyclalkylamino;
- (l) optionally substituted heterocyclalkylcarbonyl;
- (s)  $-\text{Y}-(\text{alkylene})-\text{R}^9$  where  $\text{Y}$  is a single bond,  $-\text{O}-$  or  $-\text{NH}-$  and  $\text{R}^9$  is optionally substituted heteroaryl,  $-\text{CONR}^{12}\text{R}^{13}$ ,  $\text{SO}_2\text{R}^{14}$ ,  $-\text{SO}_2\text{NR}^{15}\text{R}^{16}$ ,  $-\text{NHSO}_2\text{R}^{17}$  or  $-\text{NHSO}_2\text{NR}^{18}\text{R}^{19}$  where  $\text{R}^{12}$ ,  $\text{R}^{13}$ ,  $\text{R}^{14}$ ,  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{17}$ ,  $\text{R}^{18}$  and  $\text{R}^{19}$  are independently of each other hydrogen, alkyl or heteroalkyl;
- 20 (x) cycloalkylalkyl, cycloalkylalkynyl and cycloalkylalkynyl, all optionally substituted with alkyl, halo, hydroxy or amino;
- 25 (m) arylaminoalkylene or heteroarylaminoalkylene;
- or
- (n)  $\text{Z-alkylene-NR}^{30}\text{R}^{31}$  where  $\text{Z}$  is  $-\text{NH}-$ ,  $-\text{N}(\text{alkyl})-$  or  $-\text{O}-$ , and  $\text{R}^{30}$  and  $\text{R}^{31}$  are independently of each other, hydrogen, alkyl or heteroalkyl.
- 30

5. The compound of any one of claim 1-4, wherein  $\text{R}^1$  and  $\text{R}^2$  are hydrogen and  $\text{B}$  is phenyl.

- 118 -

6. The compound of any one of claims 1-5, wherein A is phenyl.
7. The compound of any one of claims 1-6, wherein R<sup>4</sup> is hydrogen and R<sup>5</sup> is halo or alkyl.
- 5 8. The compound of claims 1-7 wherein R<sup>5</sup> is chloro, fluoro or methyl and R<sup>6</sup> is hydrogen, chloro, fluoro, methyl or methoxy.
- 10 9. The compound of any one of claims 5-8, wherein R<sup>3</sup> is optionally substituted heteroaryl.
- 15 10. The compound of claim 9, wherein R<sup>3</sup> is pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, N-oxidopyridin-2-yl, N-oxidopyridin-3-yl, N-oxidopyridin-4-yl or pyridon-2-yl, all optionally substituted.
11. The compound of claim 9 or claim 10, wherein R<sup>3</sup> is at the 3-position.
- 20 12. The compound of any one of claims 9-11, wherein R<sup>5</sup> is 4-F or 2-Me and R<sup>6</sup> is hydrogen.
13. The compound of any one of claims 1-8, wherein R<sup>3</sup> is optionally substituted phenyl.
- 25 14. The compound of claim 13, wherein R<sup>3</sup> is 3-sulfamoylphenyl, 3-methylsulfonylphenyl, 3-carboxyphenyl or 3-ethoxycarbonylphenyl.
- 30 15. The compound of claim 13 or claim 14, wherein R<sup>3</sup> is at the 3-position.
16. The compound of any one of claims 13-15, wherein R<sup>5</sup> is 4-F and R<sup>6</sup> is hydrogen.



- 119 -

17. The compound of any one of claims 1-8, wherein R<sup>3</sup> is:

- (a) heteroalkyl;
- (b) heteroalkoxy;
- (c) heteroalkylamino;
- (d) optionally substituted heterocyclalkyl;
- (e) optionally substituted heterocyclalkoxy;
- (f) optionally substituted heterocyclalkylamino;
- (g) -Y-(alkylene)-R<sup>9</sup> where Y is a single bond, -O- or -NH- and R<sup>9</sup> is optionally substituted heteroaryl, -CONR<sup>12</sup>R<sup>13</sup>, SO<sub>2</sub>R<sup>14</sup>, -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup> -NHSO<sub>2</sub>R<sup>17</sup> or -NHSO<sub>2</sub>NR<sup>18</sup>R<sup>19</sup> where R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> are independently of each other hydrogen, alkyl or heteroalkyl; or
- (h) Z-alkylene-NR<sup>30</sup>R<sup>31</sup> where Z is -NH-, -N(alkyl)- or -O-, and R<sup>30</sup> and R<sup>31</sup> are independently of each other, hydrogen, alkyl or heteroalkyl.

18. The compound of claim 17, wherein R<sup>3</sup> is heteroalkyl.

19. The compound of claim 18, wherein R<sup>3</sup> is at the 3-position and is selected from the group consisting of 2-dimethylaminoethyl, 3-dimethylaminopropyl, 4-dimethylaminobutyl, 2-dimethylaminoethylamino, 3-dimethylaminopropylamino, hydroxymethyl, 1,2-dihydroxyethyl, 3-hydroxy-3-methyl-1-butyl or 3-hydroxybutyl.

20. The compound of claim 18 or claim 19, wherein R<sup>5</sup> is 2-F and R<sup>6</sup> is 4-F.

21. The compound of claim 18 or claim 19, wherein R<sup>5</sup> is 4-F and R<sup>6</sup> is hydrogen.

22. The compound of claim 18 or claim 19, wherein R<sup>5</sup> is 2-Me and R<sup>6</sup> is hydrogen.

- 120 -

23. The compound of claim 17, wherein R<sup>3</sup> is heteroalkoxy or heteroalkylamino.

24. The compound of claim 23, wherein R<sup>3</sup> is at the 3-position and is selected from the group consisting of 3-dimethylaminopropoxy, 2-dimethylaminoethoxy, 2-hydroxyethoxy, 2,3-dihydroxypropoxy, 2-dimethylaminoethylamino and 3-dimethylaminopropylamino.

25. The compound of claim 23 or claim 24, wherein R<sup>5</sup> is 4-F or 2-Me and R<sup>6</sup> is hydrogen.

26. The compound of claim 17, wherein R<sup>3</sup> is optionally substituted heterocyclalkyl, optionally substituted heterocyclalkoxy or optionally substituted heterocyclalkylamino.

27. The compound of claim 26, wherein R<sup>3</sup> is at the 3-position and is selected from the group consisting of 3-(morpholin-4-yl)propoxy, 2-(morpholin-4-yl)ethoxy, 2-(2-oxo-pyrrolidin-1-yl)ethoxy, 3-(morpholin-4-yl)propyl, 2-(morpholin-4-yl)ethyl, 4-(morpholin-4-yl)butyl, 3-(morpholin-4-yl)propylamino, 2-(morpholin-4-yl)ethylamino, 4-hydroxypiperidinylmethyl, 2-(S,S-dioxo-thiamorpholin-4-yl)ethyl, 3-(S,S-dioxo-thiamorpholin-4-yl)propyl and N-methylpiperazinylmethyl.

28. The compound of claim 26 or claim 27, wherein R<sup>5</sup> is 4-F or 2-Me and R<sup>6</sup> is hydrogen.

29. The compound of claim 17, wherein R<sup>3</sup> is -Y-(alkylene)-R<sup>9</sup> where Y is a single bond, -O- or -NH- and R<sup>9</sup> is optionally substituted heteroaryl, -CONR<sup>12</sup>R<sup>13</sup>, SO<sub>2</sub>R<sup>14</sup>, -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup>, -NHSO<sub>2</sub>R<sup>17</sup> or -NHSO<sub>2</sub>NR<sup>18</sup>R<sup>19</sup> where R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> are independently of each other hydrogen, alkyl or heteroalkyl.

- 121 -

30. The compound of claim 29, wherein Y is a single bond and R<sup>9</sup> is SO<sub>2</sub>R<sup>14</sup> or -SO<sub>2</sub>NR<sup>15</sup>R<sup>16</sup>.

31. The compound of claim 29 or claim 30, wherein R<sup>3</sup> is methylsulfonylethyl or sulfamoylethyl.

32. The compound of any one of claims 29-31, wherein R<sup>5</sup> is 4-F or 2-Me and R<sup>6</sup> is hydrogen.

33. A compound selected from the group consisting of:

5-amino-1-(4-fluorophenyl)-4-[3-(2-morpholin-4-ylethoxy)-benzoyl]pyrazole,

5-amino-1-(2,4-difluorophenyl)-4-[3-(3-morpholin-4-ylpropyl)benzoyl]pyrazole,

5-amino-4-(3-aminobenzoyl)-1-(4-fluorophenyl)pyrazole,

5-amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylpropyl)-benzoyl]pyrazole,

5-amino-4-[3-(2-aminosulfonylethenyl)benzoyl]-1-(4-fluorophenyl)pyrazole,

5-amino-4-(3-acetylaminobenzoyl)-1-phenylpyrazole,

5-amino-4-[3-(2-aminoethyl)benzoyl]-1-(4-fluorophenyl)pyrazole,

5-amino-1-(4-fluorophenyl)-4-[3-(3-morpholin-4-ylpropylamino)-benzoyl]pyrazole,

5-amino-4-[3-(2-aminosulfonylethyl)benzoyl]-1-(4-fluorophenyl)-pyrazole,

5-amino-1-(4-fluorophenyl)-4-(3-pyridin-3-ylbenzoyl)pyrazole,

and

34. A compound selected from the group consisting of:

5-amino-1-(2-methylphenyl)-4-[3-pyridin-3-yl)benzoyl]pyrazole,

5-amino-1-(2-methylphenyl)-4-[3-(N-oxidopyridin-3-yl)benzoyl]-pyrazole,

5-amino-4-[3-(2,3-dihydroxypropoxy)benzoyl]-1-(4-fluorophenyl)-pyrazole,

- 122 -

5-amino-4-[3-(1,2-dihydroxyethyl)benzoyl]-1-(4-fluorophenyl)-  
pyrazole,

5-amino-1-(4-fluorophenyl)-4-[3-(sulfamoylbenzoyl)pyrazole,

5-amino-1-(4-fluorophenyl)-4-[3-(3-hydroxy-3-methylbutyl)-  
benzoyl]pyrazole,


5-amino-1-(4-fluorophenyl)-4-[3-(2-(1-hydroxycyclopentyl)ethyl)-  
benzoyl]pyrazole,

5-amino-4-[3-(2-methylsulfonyl)ethyl]benzoyl]-1-(4-fluorophenyl)-  
pyrazole, and


5-amino-1-(2,4-difluorophenyl)-4-[3-(2-hydroxyethylsulfonyl)-  
benzoyl]pyrazole.

35. A process for preparing a compound of Formula (I) selected from  
compounds of Claim 1, which process comprises:

(i) reacting a 2-keto-3-phenylaminoacrylonitrile of Formula 1:




with a hydrazine of Formula 2:



where R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are as defined in Claim 1 to provide a  
compound of Formula (I) where R<sup>1</sup> is hydrogen; or


(ii) reacting a 2-keto-3-phenylaminoacrylonitrile of formula 3:



where Z is either hydroxy, nitro or halo group and R<sup>4</sup> are as defined  
in Claim 1 with a hydrazine of formula 2 to provide a compound of formula

4:


- 123 -



followed by conversion of the Z group to the desired R<sup>3</sup> group to provide a compound of Formula (I) where R<sup>1</sup> is hydrogen;

- (iii) optionally modifying any of the R<sup>1</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> or R<sup>6</sup> groups;
- (iv) optionally converting the compound of Formula (I) prepared in Steps (i), (ii) or (iii) above, to the corresponding acid addition salt by treatment with an acid;
- (v) optionally converting the compound of Formula (I) prepared in Steps (i), (ii) or (iii) above, to the corresponding free base by treatment with a base; and
- (vii) optionally separating a mixture of stereoisomers of a compound of Formula (I) prepared in Steps (i) - (v) above, to give a single stereoisomer.

36. A process for preparing a compound of Formula (I) selected from compounds of Claim 1, which process comprises reacting a compound of Formula 5:



- where R<sup>5</sup> and R<sup>6</sup> are as in claim 1 and L is a leaving group under organometallic displacement reaction conditions


- 124 -



with an organometallic reagent of formula  $\text{R}^3\text{A}(\text{R}^4)\text{M}^+$  where  $\text{R}^3$  and  $\text{R}^4$  are as in claim 1 and M is a metallic moiety to provide a compound of Formula (I) where  $\text{R}^1$  is hydrogen;

- (ii) optionally modifying any of the  $\text{R}^1$ ,  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$  or  $\text{R}^6$  groups;
  - (iii) optionally converting the compound of Formula (I) prepared in Steps (i) or (ii) above, to the corresponding acid addition salt by treatment with an acid;
  - (iv) optionally converting the compound of Formula (I) prepared in Steps (i) or (ii) above, to the corresponding free base by treatment with a base; and
- optionally separating a mixture of stereoisomers of a compound of Formula (I) prepared in Steps (i) or (iv) above, to give a single stereoisomer.

37. A compound of formula



wherein Z is either hydroxy, nitro or halo group and A, B,  $\text{R}^4$ ,  $\text{R}^5$  and  $\text{R}^6$  are as defined in claim 1.

38. A compound according to any one of claims 1-34 for use as a medicament.

39. A medicament containing a compound according to any one of claims 1-34 and a pharmaceutically acceptable excipient.

40. A medicament of claim 39 for the control or prevention of inflammatory diseases.

- 125 -

5 41. A process for the manufacture of medicaments which process  
comprises bringing a compound according to any one of claims 1-  
34 together with a pharmaceutically acceptable excipient and  
bringing the mixture into a galenical administration form.

42. The use of a compound of any one of claims 1-34 in the treatment  
or prevention of inflammatory diseases.

10 43. The use of a compound of any one of claims 1-34, for the  
manufacture of a medicament for the treatment and prophylaxis  
of inflammatory diseases.

15 44. A compound according to any one of claims 1-34 whenever  
prepared according to a process of claim 35 or claim 36.

45. The novel compounds, intermediates, processes, formulations as  
hereinbefore described.

20

\* ... \* ... \* ... \* ... \*

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/02879

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D231/38 A61K31/415 C07D401/10 C07D403/12 C07D401/12  
A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TUPPER, DAVID E. ET AL: "Steric and electronic control in the addition of hydrazine and phenylhydrazine to .alpha.-'(dimethylamino)methylene!-.beta.- oxoarylpropanenitriles" SYNTHESIS (1997), (3), 337-341 , XP002115275 compounds 5b,5e ----- -/--	37



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&amp;" document member of the same patent family

Date of the actual completion of the international search

15 September 1999

Date of mailing of the international search report

05/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/02879

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NISHIWAKI, TAROZAEMON ET AL: "Heterocyclic chemistry. XIX. Synthesis of 4-aryl-1-arylpyrazoles from .alpha.-aryl-.beta.-anilinoacrylonitr iles and photochemistry of 4-carbonyl substituted pyrazoles" J. CHEM. SOC., PERKIN TRANS. 1 (1974), (15), 1871-5 ; XP002115276 p-chlorophenyl-substituted compounds of formula (2) page 1871 -page 1872 -----	37
A	WO 94 13643 A (PFIZER ; FARACI WILLIAM STEPHEN (US); WELCH WILLARD MCKOWAN JR (US)) 23 June 1994 (1994-06-23) claim 1; examples -----	1, 39

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/02879

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 42  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim 42  
is directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compounds.
2. ☒ Claims Nos.: 45  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 45

It is not clear to which compounds, intermediates, processes and formulations, claim 45 refers. The present application therefore fails to comply with the clarity requirements of Article 84 EPC to such an extent, that a meaningful search for claim 45 is impossible.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/02879

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9413643 A	23-06-1994	AT 175961 T	15-02-1999
		AU 690527 B	30-04-1998
		AU 5454894 A	04-07-1994
		AU 7843198 A	01-10-1998
		CA 2150483 A	23-06-1994
		CN 1092768 A	28-09-1994
		CZ 9501585 A	17-01-1996
		DE 69323212 D	04-03-1999
		DE 69323212 T	27-05-1999
		EG 20397 A	28-02-1999
		EP 0674624 A	04-10-1995
		ES 2126661 T	01-04-1999
		FI 935674 A	18-06-1994
		HU 67457 A	28-04-1995
		IL 107946 A	24-09-1998
		JP 2862374 B	03-03-1999
		JP 7509725 T	26-10-1995
		MX 9308042 A	31-08-1994
		NO 952395 A	16-08-1995
		NO 985494 A	16-08-1995
		NZ 257770 A	24-02-1997
		PL 309356 A	02-10-1995
		US 5712303 A	27-01-1998
		ZA 9309404 A	15-06-1995